



Whole-blood fatty acids are associated with executive function in Tanzanian children aged 4–6 years: a cross-sectional study

Theresia Jumbe^{1,3}, Sarah S. Comstock¹, William S. Harris², Joyce Kinabo³, Matthew B. Pontifex⁴ and Jenifer I. Fenton^{1*}

¹Department of Food Science and Human Nutrition, Michigan State University, MI 48824, USA

²Sanford School of Medicine, University of South Dakota and OmegaQuant Analytics, LLC, Sioux Falls, SD 57106, USA

³Sokoine University of Agriculture, Department of Food Technology, Nutrition and Consumer Sciences, Morogoro, Tanzania

⁴Department of Kinesiology, Michigan State University, MI 48824, USA

(Submitted 23 November 2015 – Final revision received 19 August 2016 – Accepted 3 September 2016 – First published online 21 October 2016)

Abstract

Essential fatty acids (EFA) are PUFA that are metabolised to long-chain PUFA and are important for brain development and cognitive function. The objective of this study was to determine the association between whole-blood EFA and cognitive function in Tanzanian children. A total of 325 2–6-year-old children attempted the dimensional change card sort (DCCS) tasks to assess executive function. Blood samples were collected for fatty acid (FA) analysis by GC. Associations between executive function and FA levels were assessed by regression. Among the 130 4–6-year-old children who attempted the DCCS tasks, whole-blood levels of linoleic acid were positively associated with executive function, whereas whole-blood levels of α -linolenic acid and nervonic acid were inversely associated with executive function. A full model including all twenty-five FA explained 38% of the variation in executive function, whereas a reduced model including only the EFA (α -linolenic acid and linoleic acid), DHA and EPA explained 25% of the variation in executive function. Children who had sufficient whole-blood levels of EFA were 3.8 times more likely to successfully complete all DCCS tasks compared with children with insufficient EFA. These results suggest that whole-blood FA levels are associated with cognitive abilities. Intervention trials that include assessment of whole-blood FA levels are required to determine the relationships between intake, blood levels and executive function in Tanzanian children.

Key words: Cognition: Executive function: Fatty acids: Lipids: Essential fatty acids: Brain development: Tanzania

Long-chain PUFA (LCPUFA) accumulate in the fetus during pregnancy and during early childhood⁽¹⁾. These PUFA are concentrated in the central nervous system⁽²⁾. Essential fatty acids (EFA) of both *n-6* and *n-3* fatty acid (FA) families and their LCPUFA metabolites play a significant role in neuronal growth and differentiation of cells and have been associated with cognitive abilities of children^(2–4). Poor PUFA status may affect brain development as well as the cognitive abilities of children⁽⁴⁾. Brain development continues through childhood and early adolescence, with cerebral volume reaching 95% of its peak by 6 years of age and reaching its maximum between 10 and 15 years of age⁽⁵⁾. Thus, LCPUFA should be included in the diets of infants and children to ensure optimal brain development^(6–8).

In most developing countries, a large proportion of the population cannot afford diets rich in animal foods^(4,9), and lack of foods from animal sources may lead to PUFA deficiency⁽⁴⁾. Several PUFA supplementation studies conducted in young

children (< 2 years) have demonstrated that children fed foods/milk fortified with α -linolenic acid (ALA) or its metabolites EPA and DHA alone or together with other micronutrients enhanced cognitive development⁽⁹⁾. These studies suggest that high intake of LCPUFA may improve cognitive abilities later in life (i.e. after 2 years of age)^(9,10). Sheppard & Cheatham⁽¹⁰⁾ concluded that LCPUFA influence the cognitive development of children, especially with regard to planning and memory processing. Previous studies have utilised food intake data or supplementation programmes to estimate FA status of children⁽¹¹⁾. To our knowledge, a few of these studies have directly measured whole-blood FA status as it relates to cognitive development in children between 4 and 6 years of age.

Executive function, the conscious control of thoughts and actions, develops between the ages of 2 and 10 years⁽¹²⁾. Executive function involves inhibition, working memory and task switching⁽¹²⁾ and is controlled by the frontal and temporal lobes of the brain⁽¹⁰⁾. These two regions of the brain continue to

Abbreviations: ALA, α -linolenic acid; BAZ, BMI-for-age z-score; EFA, essential fatty acid(s); EFAD, essential fatty acid deficiency/deficient; FA, fatty acid(s); LA, linoleic acid; LCPUFA, long-chain PUFA; T:T, triene:tetraene.

* **Corresponding author:** J. I. Fenton, email imigjeni@msu.edu

develop after the 2nd year of life and contain high amounts of arachidonic acid (AA) and DHA⁽¹³⁾. A method commonly used to assess executive function in young children is the dimensional change card sorting (DCCS) task⁽¹²⁾. Therefore, we used the DCCS task to assess cognitive function in this population of young children.

In addition, although FA are widely understood to affect growth and cognition, the associations between blood levels of specific FA and these health outcomes are infrequently reported. In this study, we assess the association between FA status and executive function in Tanzanian children using a DCCS test with culturally modified colours and images⁽¹²⁾. We hypothesised that whole-blood levels of EPA, DHA and both EFA (ALA and linoleic acid) would be positively associated with performance on the DCCS tasks.

Methods

Study site

The present study was conducted in Rudewa Mbuyuni village in Kilosa District, Morogoro, Tanzania. Conditions prevailing in the village have previously been described⁽¹⁴⁾. Children in the village begin attending primary school by about 7 years of age. At present, there are no preschool programmes in the village. The Ministry of Health in Tanzania requires that all children <5 years of age visit a growth-monitoring clinic every month, receive vaccinations according to a published schedule as well as receive vitamin A drops twice a year (June, December).

Subjects and ethics approval

All children in the village were invited to participate in this study. From December 2013 to August 2014, 335 apparently healthy children between 2 and 6 years of age were enrolled in this cross-sectional study. All participants and their mothers/caregivers verbally consented to participate in the study. The National Medical Research Board (NIMR) of Tanzania (NIMR/HQ/R.8a/Vol. IX/1189) and the Michigan State University (USA) Human Research Protection Program (IRB no. 13-700) approved the study. These children were also enrolled in an International Development Research Center (Ottawa, ON, Canada)-funded research study entitled Food Security, Adequate Care and Environment: Eco-Nutrition Guidelines for Community Action on Climate Change. Some results from that study have been previously published^(14,15). A total of ten children refused to take the test at initial contact and withdrew from the study. Swahili was used as the language of communication throughout the study.

Anthropometric measurements

Height was measured to the nearest 0.1 cm with a stadiometer (Shorr Productions; Perspective Enterprises), and weight was measured using a digital bathroom scale to the nearest 0.1 kg (SECA; Vogel & Haiké). The average of two measurements was used. Date of birth was recorded from the reproductive and child health clinic (RCH) card, and mother's recall was used for those without a RCH card. Sex of the child was recorded.

Data on weight, height, date of birth and sex were entered into World Health Organization Anthro⁽¹⁶⁾ and World Health Organization AnthroPlus⁽¹⁷⁾ to calculate z-scores.

Whole-blood assessments

A capillary blood sample was obtained from the middle finger for malaria rapid test (Premier Medical Co. Ltd) and measurement of Hb concentrations using a HemoCue photometer (HemoCue AB); samples were dropped onto dried blood spot (DBS) cards pre-treated with an antioxidant cocktail⁽¹⁸⁾. DBS cards were stored in a dry, dark, cool environment and shipped to OmegaQuant in USA for FA analysis within 14 d of sample collection. DBS cards were punched and combined with a derivatising reagent (boron trifluoride in methanol (14%), toluene and methanol (35:30:35 parts)), shaken and heated at 100°C for 45 min. After cooling, forty parts of both hexane and distilled water were added and briefly vortexed. FA methyl esters were analysed using a GC as previously described⁽¹⁹⁾. The stability of FA from blood samples collected, stored and transported in this manner has been documented in previous studies from our laboratory^(20,21) and by others^(22,23). Whole-blood FA proportions are expressed as a percentage of total identified FA. The following FA were assessed: myristic, palmitic, palmitelaidic, palmitoleic, stearic, elaidic, oleic, linoelaidic, linoleic, arachidic, γ -linolenic, eicosanoic, ALA, eicosadienoic, behenic, mead, dihomo- γ -linolenic, AA, lignoceric, EPA, nervonic, docosatetraenoic, DHA, docosapentaenoic *n*-6 and docosapentaenoic *n*-3. The triene:tetraene (T:T) ratio is the ratio of mead acid:AA. Historically, a T:T ratio > 0.02 in plasma samples defines essential FA deficiency^(24,25). Therefore, a T:T ratio > 0.02 was used to define insufficient levels of EFA^(26,24).

Cognitive assessment: dimensional change card sort

The DCCS^(12,27) is conceptually simplistic in that it requires the child to sort a series of bivalent cards (online Supplementary Fig. S1) on the basis of one of two instructed dimensions (i.e. colour or shape). Following sorting of an initial series of eight cards based upon colour, the children were instructed to switch the categorisation dimension and sort another series of eight cards on the basis of shape.

Previous studies have demonstrated that children younger than 3 years of age can complete the pre-switch series⁽²⁸⁾, but the dimensional change requires engagement of executive function in order to inhibit the previous rule set to execute the correct sorting behaviour^(12,29). Indeed, children with poor executive function exhibit a tendency to perseverate during the post-switch series by continuing to sort the cards by the first dimension despite being able to verbally express the new sorting rules. A critical limitation of the traditional card sort task, however, is the relatively narrow age range in which it can be utilised, with Rennie *et al.*⁽²⁹⁾ observing that children are unable to successfully complete the post-switch sorting series until about 4–5 years of age. Accordingly, a modified variant of the DCCS task was used in order to increase the number of participants who could perform the task⁽²⁸⁾. As young children and those with poor executive function appear to have greater attentional inertia – manifesting with increased difficulty

separating features of an object⁽³⁰⁾ – separating the sorting attributes into differentiable objects reduces, but does not remove, the inhibitory demands required to complete the switch⁽²⁸⁾. Therefore, participants completed three versions of the DCCS, which progressively increased the executive function requirements by increasing the overlap between the shape and colour stimulus on the cards (i.e. no overlap, partial overlap, full overlap). The task was modified to use images and colours familiar to the participating children. The mother or caregiver was present during the test to observe the process and allow the child to feel comfortable and confident.

Each version of the DCCS (no overlap, partial overlap, full overlap) had a pre- and post-switch phase. For a child to pass any phase, she or he needed to obtain six correct responses out of eight. If fewer than six correct responses were made in the pre-switch phase, the post-switch phase of that version was not scored. The child was then asked to continue with the next version of the DCCS. Consistent with two dominant approaches to scoring the DCCS, task performance was summarised using the following: (1) highest test passed and (2) total passes. Scoring for (1) 'highest test passed' was ordered. A child scored '0' if he or she was unable to pass any post-switch phase, '1' if the child passed the no overlap DCCS post-switch task and the partial overlap post-switch, and '2' if the child passed the full overlap post-switch task. Scoring for (2) 'total passes' was not ordered. It was based on the total number of post-switch phases passed and ranged from 0 to 3.

Data reduction and statistical analyses

An *a priori* power analysis was conducted using the results of previous investigations observing an association between nutritional supplementation and performance on the proposed tasks⁽³¹⁾. Assuming a conservative effect size ($f^2=0.1$),

a two-sided α of 0.05 and a β of 0.20 (i.e. 80% power), a sample of eighty-one participants was estimated to provide adequate power. As more than half of the children <48 months of age failed to pass any DCCS task (Fig. 1), we excluded them from our analyses. The analyses presented in this study are from the 130 children older than 48 months. A *post hoc* analysis of the statistical power using the method of Cohen⁽³²⁾ was conducted. Using the data obtained from the analysis of linoleic acid, with 130 subjects, and α set to 0.05, we had 92% power to detect an R^2 of 0.08. This was adjusted for two additional independent variables (Hb and malaria) with an R^2 of 0.07.

Basic descriptive analyses were conducted to obtain means and frequencies. Means between groups (i.e. the total study population *v.* the 130 kids \geq 48 months of age; those who did not pass any DCCS task *v.* those who passed at least one DCCS task) were compared using *t* tests (for continuous data) or χ^2 analysis (for proportions), as appropriate.

As high collinearity among the FA caused tolerance levels < 0.1 and gave variance inflation factors >10 when all the FA were entered into a single model, single linear regressions were used to analyse the association between blood FA levels and executive function. Models for linear regression included the FA of interest, Hb levels and malaria status. We included Hb concentrations in our model because it is a known significant predictor of cognitive abilities in our population⁽³³⁾. None of the children exhibited symptoms of active malaria infection. Positive malaria status indicated subclinical infection and was included as a covariate because it was positively associated with DCCS performance. To correct for multiple testing, PROC MULTTEST with the false discovery rate (FDR) option was applied. The FDR option uses the linear step-up adjustment described by Benjamini & Hochberg⁽³⁴⁾. SAS version 9.4 was used for these statistical analyses.

Factor analysis was used to reduce the number of FA variables using Proc Factor in SAS version 9.4. This enabled




Fig. 1. Percentage of children able to pass each stage of the dimensional change card sort test. ■, Did not pass; □, passed no overlap; ▨, passed partial overlap; ▩, passed full overlap.

correlated variables to be assessed simultaneously. A linear transformation was performed to enable interpretation. In this case, varimax rotation was performed; three factors were retained as determined by eigenvalues > 2. For each of the three factors, FA with rotated factor loadings > 0.2 constitute that factor. The procedure assigns each person a score for each of the three factors that emerged from the data. Multiple linear regression using these factor scores as predictors was used to determine the associations between these factors and performance on the DCCS tasks. Models for these linear regressions included the three factor scores, age, sex, BMI-for-age z-scores (BAZ), Hb levels and malaria status. To determine associations between EFA status and executive function, we conducted polytomous logistic regressions for categorical variables using SAS version 9.4. In all cases, a *P* value < 0.05 was used to define statistical significance.

Results

This study enrolled 335 children between 2 and 6 years of age. Of these, 325 attempted the DCCS tasks and ten refused to attempt the DCCS. Basic information about the study participants are presented in Table 1. Less than half of the children between 24 and 48 months successfully performed any DCCS task (Fig. 1). Of the 130 children ≥ 48 months who attempted the DCCS tasks, 38% passed the full overlap task, 18% passed the partial overlap task, 13% passed the no overlap task and 31% failed to successfully complete any of the tasks. Children ≥ 48 months who successfully completed at least one DCCS task were older, taller and more likely to test positive for malaria than those children who could not complete any DCCS tasks (Table 2).

FA levels in whole blood from the 130 children whose data were analysed are presented in Table 3. The mean linoleic acid level was 17.6 (SD 2.7), as a percent of whole-blood FA, whereas the mean ALA level was 0.4 (SD 0.2). The mean duration of breast-feeding among children ≥ 48 months was 22.8 (SD 4.4) months. Breast-feeding duration was similar between children who failed to complete any DCCS task and those who successfully completed DCCS tasks (Table 2). In this population, breast-feeding duration closely matched the WHO recommendation to breast-feed up to 24 months⁽³⁵⁾.

Table 1. Participant characteristics (Mean values and standard deviations; numbers and percentages)

	Overall (<i>n</i> 325)		Children ≥ 48 months (<i>n</i> 130)		<i>P</i>
	Mean	SD	Mean	SD	
Age (months)	45.33	14.70	61.04	7.43	<0.01
Sex (male)					0.75
<i>n</i>	149		59		
%	46.5		45.4		
Height (cm)	94.48	9.56	103.14	6.29	<0.01
Weight (kg)	13.92	2.61	16.04	2.07	<0.01
Hb (g/l)	102.9	14.6	105.8	14.5	<0.01
Malaria (%)	16.3		20.0		0.04
Duration of breast-feeding (months) (<i>n</i> 312)	22.42	4.06	22.75	4.36	0.25

Regression results between the selected FA and the ordered assessment (the highest test passed) and the non-ordered assessment (total passes) of DCCS performance are shown in Table 4. A significant inverse association was observed between DCCS performance and ALA and nervonic acid. Linoleic acid was positively associated with DCCS performance. These three FA were tested in a multiple linear regression model that included the following confounders: malaria infection status, Hb concentration, age, sex and BAZ. For both non-ordered and ordered assessments of DCCS performance, the model was significant (*P* < 0.0001). For the non-ordered assessment, this

Table 2. Characteristics of children ≥ 48 months stratified by dimensional change card sort performance (Mean values and standard deviations; numbers and percentages)

	Failing (<i>n</i> 42)		Passing (<i>n</i> 88)		<i>P</i>
	Mean	SD	Mean	SD	
Age (months)	59.14	7.30	61.95	7.37	0.04
Sex (male)					0.47
<i>n</i>	21		38		
%	50.0		43.2		
Height (cm)	101.37	5.82	103.99	6.37	0.03
Weight (kg)	15.68	1.82	16.22	2.17	0.16
Hb (g/l)	106.3	14.5	105.6	14.7	0.79
Malaria (%)					0.02
%	9.52		27.3		
Duration of breast-feeding (months) (<i>n</i> 126)	22.00	5.36	23.08	3.82	0.26
BAZ	-0.06	0.75	-0.29	0.90	0.14
WAZ	-1.11	0.97	-1.09	0.82	0.86

BAZ, BMI-for-age z-score; WAZ, weight-for-age z-score.

Table 3. Whole blood fatty acid proportions in Tanzanian children ≥ 48 months of age (*n* 130) (Mean values and standard deviations)

	Mean	SD	Range
Myristic	0.80	0.45	0.21–2.77
Palmitic	25.36	1.98	21.11–31.50
Oleic	21.47	3.15	14.30–33.57
Linoleic	17.58	2.74	12.13–26.78
α-Linolenic	0.41	0.19	0.12–1.32
Mead	0.14	0.06	0.02–0.29
Arachidonic	10.04	1.65	4.03–14.02
EPA	0.43	0.18	0.07–1.03
DHA	2.94	0.76	1.04–5.01
Palmitelaidic	0.09	0.07	0.00–0.44
Palmitoleic	1.53	0.64	0.46–3.81
Stearic	10.57	1.04	6.88–12.91
Elaidic	0.22	0.31	0.08–3.54
Linoelaidic	0.33	0.11	0.15–1.14
Arachidic	0.26	0.06	0.13–0.46
γ-Linolenic	0.31	0.13	0.09–0.72
Eicosenoic	0.25	0.08	0.13–0.53
Eicosadienoic	0.27	0.09	0.13–0.61
Behenic	0.62	0.20	0.15–1.19
Dihomo-γ-linolenic	1.77	0.42	1.01–3.78
Lignoceric	0.82	0.39	0.17–1.93
Nervonic	0.70	0.32	0.11–1.64
Docosatetraenoic	1.35	0.32	0.48–2.04
Docosapentaenoic (<i>n</i> -6)	0.78	0.16	0.39–1.20
Docosapentaenoic (<i>n</i> -3)	0.95	0.26	0.47–2.04

Table 4. Regression† results for the two methods of scoring the dimensional change card sort and selected fatty acids (FA)

FA	Regression results for total passes				Regression results for highest test passed			
	Parameter estimate	Standardised parameter estimate	Raw <i>P</i>	FDR <i>P</i>	Parameter estimate	Standardised parameter estimate	Raw <i>P</i>	FDR <i>P</i>
Arachidonic	0.09	0.12	0.18	0.30	0.08	0.11	0.23	0.37
Stearic	0.24	0.20	0.03	0.11	0.27	0.22	0.02	0.08
Docosatetraenoic	-0.11	-0.03	0.75	0.84	-0.16	-0.04	0.64	0.78
Docosapentaenoic (<i>n</i> -3)	-0.51	-0.10	0.22	0.36	-0.44	-0.09	0.30	0.45
Dihomo- γ -linolenic	0.30	0.10	0.24	0.37	0.43	0.14	0.10	0.23
Docosapentaenoic (<i>n</i> -6)	-0.04	0.00	0.96	0.96	-0.25	-0.03	0.72	0.78
EPA	-0.99	-0.14	0.11	0.28	-0.75	-0.10	0.23	0.37
DHA	0.27	0.17	0.05	0.18	0.27	0.16	0.07	0.18
Eicosadienoic	-0.76	-0.05	0.54	0.70	-0.64	-0.04	0.62	0.78
Behenic	-0.76	-0.12	0.16	0.30	-0.96	-0.15	0.09	0.23
Arachidic	-2.52	-0.12	0.15	0.28	-2.95	-0.14	0.10	0.23
Lignoceric	-0.60	-0.19	0.03	0.11	-0.67	-0.20	0.02	0.08
Nervonic*	-0.91*	-0.23*	0.01*	0.04*	-1.02*	-0.25*	<0.01*	0.02*
Palmitelaidic	1.37	0.08	0.39	0.56	0.43	0.02	0.79	0.81
Elaidic	0.21	0.05	0.54	0.70	0.14	0.03	0.70	0.78
α -Linolenic*	-1.61*	-0.25*	<0.01*	0.04*	-1.53*	-0.23*	0.01*	0.05*
Eicosenoic	-0.63	-0.04	0.64	0.75	-1.00	-0.06	0.46	0.66
Palmitoleic	-0.25	-0.13	0.14	0.28	-0.26	-0.13	0.13	0.27
Mead	-1.03	-0.05	0.59	0.73	-1.16	-0.05	0.56	0.75
Palmitic	-0.14	-0.22	0.01	0.07	-0.13	-0.19	0.03	0.11
γ -Linolenic	-0.06	-0.01	0.94	0.96	-0.35	-0.03	0.69	0.78
Myristic	-0.35	-0.12	0.15	0.28	-0.37	-0.13	0.14	0.27
Oleic	-0.06	-0.15	0.09	0.26	-0.07	-0.17	0.06	0.18
Linoelaidic	-0.07	-0.01	0.94	0.96	-0.24	-0.02	0.81	0.81
Linoleic*	0.13*	0.29*	<0.01*	0.02*	0.14*	0.30*	<0.01*	0.02*

FDR, false discovery rate.

*All significant associations ($P < 0.05$).

† Model: total passes = FA + malaria status + Hb concentration. Highest test passed = FA + malaria status + Hb concentration.

model explained 24% of the variation (r^2 0.24; adjusted r^2 0.19). For the ordered assessment, this model explained 25% of the variation (r^2 0.25; adjusted r^2 0.20). To test the hypothesis that whole-blood levels of EPA, DHA and both EFA (ALA and linoleic acid) would be positively associated with performance on the DCCS tasks, multiple linear regression using EPA, DHA, ALA, linoleic acid, malaria, Hb, age, sex and BAZ was conducted. For both non-ordered and ordered assessments of DCCS performance, the model was significant ($P \leq 0.0002$). For the non-ordered assessment, this model explained 26% of the variation (r^2 0.26; adjusted r^2 0.21). For the ordered assessment, this model explained 24% of the variation (r^2 0.24; adjusted r^2 0.18). DHA ($p \leq 0.01$), ALA ($P < 0.05$), linoleic acid ($p \leq 0.01$) and malaria ($p \leq 0.01$) were significant contributors to the model in both non-ordered and ordered assessments. DHA, linoleic acid and malaria were positively associated with performance on the DCCS tasks, whereas ALA was inversely associated with performance on the DCCS tasks. It is notable that a full model including all twenty-five single FA as well as Hb concentrations, malaria status, age, sex and BAZ was significant ($P = 0.003$ for total passes and $P = 0.004$ for highest test passed) and explained about 38% of the variance (r^2 0.38; adjusted r^2 0.21 for total passes and adjusted r^2 0.19 for highest test passed). However, the effects of independent FA and the covariates could not be determined because of high levels of collinearity causing poor tolerance and variance inflation in the model.

To bypass the problems with collinearity, factor analysis was conducted. When factor analysis was used to determine how combinations of the FA might be associated with

Table 5. Factor loading matrix for fatty acids (FA) in the whole blood of Tanzanian children ≥ 48 months of age

	Factor 1	Factor 2	Factor 3
Arachidonic	0.8550	0.0602	-0.2799
Stearic	0.7802	-0.0516	-0.2820
Docosatetraenoic	0.6861	0.1677	-0.0039
Docosapentaenoic (<i>n</i> -3)	0.6335	0.1461	0.1777
Dihomo- γ -linolenic	0.6059	-0.1102	0.0670
Docosapentaenoic (<i>n</i> -6)	0.6057	0.1607	0.0215
EPA	0.4746	0.2805	0.3291
DHA	0.4477	0.1974	-0.4181
Eicosadienoic	0.3589	0.2227	-0.1069
Behenic	0.3334	0.8293	-0.1706
Arachidic	0.1079	0.8839	-0.1059
Lignoceric	0.2361	0.8590	-0.1079
Nervonic	0.2423	0.8486	-0.1149
Palmitelaidic	0.0642	0.4690	-0.2016
Elaidic	-0.2726	0.4142	-0.2687
α -Linolenic	0.0373	0.3622	0.1246
Eicosenoic	-0.0153	0.2200	0.0578
Palmitoleic	-0.1676	-0.2225	0.7625
Mead	0.1084	-0.1114	0.7158
Palmitic	-0.4937	-0.2078	0.6712
γ -Linolenic	0.1296	-0.0768	0.5274
Myristic	-0.1163	-0.1058	0.4231
Oleic	-0.7860	-0.0563	0.3633
Linoelaidic	-0.1042	0.1126	0.2933
Linoleic	0.0105	-0.2277	-0.7862

performance on the DCCS tasks, three factors emerged. The factor loading matrix is shown in Table 5. Multiple linear regression using these three factors ($P < 0.0001$) demonstrated

Table 6. Regression† results for the two methods of scoring the dimensional change card sort and fatty acid (FA) factors

	Regression results for total passes			Regression results for highest test passed		
	Parameter estimate	Standardised parameter estimate	<i>P</i>	Parameter estimate	Standardised parameter estimate	<i>P</i>
Factor 1	0.07	0.06	0.51	0.09	0.07	0.39
Factor 2	-0.34*	-0.28*	<0.01*	-0.39*	-0.30*	<0.01*
Factor 3	-0.38*	-0.30*	<0.01*	-0.38*	-0.30*	<0.01*
Hb	0.02	0.03	0.76	-0.00	-0.00	0.98
Malaria	1.04*	0.34*	<0.01*	1.03*	0.33*	<0.01*
Age	0.04*	0.21*	0.01*	0.03*	0.17*	0.04*
Sex	-0.10	-0.04	0.63	-0.15	-0.06	0.49
BMI-for-age	0.09	0.06	0.44	0.07	0.04	0.60

*All significant associations ($P < 0.05$).

† Model: total passes = factor 1 + factor 2 + factor 3 + malaria status + Hb concentration + age + sex + BMI-for-age z-score; model P value, $P < 0.0001$; r^2 0.26; adjusted r^2 0.21. Highest test passed = factor 1 + factor 2 + factor 3 + malaria status + Hb concentration + age + sex + BMI-for-age z-score; model P value, $P < 0.0001$; r^2 0.26; adjusted r^2 0.21.

that factor 2 ($P < 0.01$) and factor 3 ($P < 0.01$) were significantly inversely associated with performance on the DCCS tasks (Table 6). When combined with Hb concentrations, malaria status, age, sex and BAZ, these parameters explained 26% of the variance (r^2 0.26; adjusted r^2 0.21 for both ordered and non-ordered assessments) in the performance on the DCCS tasks.

Polytomous logistic regression analyses demonstrated that children with low EFA levels (T:T ratio > 0.02) tended to perform more poorly on DCCS tasks than children with high EFA levels (T:T ratio ≤ 0.02). These models included malaria status and Hb levels as co-variables. For the non-ordered assessment of DCCS performance (total passes), children with higher levels of EFA were seven times more likely to successfully complete all three post-switch DCCS tasks than children with lower levels of EFA (OR 6.9; 95% CI 1.4, 35.3; $P = 0.02$). The overall model P value was 0.13 when Hb and malaria were included and 0.09 when they were not included. This was also true for the ordered assessment of DCCS performance, where children with higher levels of EFA were four times more likely to successfully complete the full overlap post-switch DCCS task than children with lower levels of EFA (OR 3.8; 95% CI 1.05, 13.9; $P = 0.04$). The overall model P value was 0.13 when Hb and malaria were included and 0.09 when they were not included. The inclusion of Hb and malaria in the models did not affect the OR, CI or P values for the EFA levels comparisons.

Discussion

The hypothesis that children with higher whole-blood levels of EFA would be more likely to successfully complete the DCCS tasks was partially supported. When whole-blood FA levels were analysed individually, children with higher levels of linoleic acid exhibited better executive function. However, children with higher levels of ALA exhibited poorer executive function. When analysed individually, neither DHA nor EPA was associated with executive function. In a model that simultaneously included the parameters EPA, DHA, ALA, linoleic acid, malaria status, Hb concentration, age, sex and BAZ, DHA, linoleic acid and malaria status were positively associated with executive function, and ALA was inversely associated with executive function. Furthermore, children with sufficient EFA

levels (T:T ratio ≤ 0.02) were four times more likely to pass the full overlap post-switch DCCS task than children with low EFA levels.

A model that included all twenty-five FA, malaria status, Hb concentration, age, sex and BAZ indicated that these parameters explained 20% (adjusted r^2) to 38% (r^2) of the variation in executive function in Tanzanian children of 4–6 years of age. This may be indicative of the importance of energy status in human brain development as the human brain uses 44–87% of resting metabolic energy during childhood⁽³⁶⁾. The brain's peak use of daily energy occurs by about 4 years of age⁽³⁶⁾. The fact that the n -6 FA linoleic acid, which is critically required for the efficient use of dietary energy content⁽³⁷⁾, is positively associated with executive function supports the idea that energy availability may be key for optimal cognitive performance. Unfortunately, total energy intake and resting metabolic rate data were not available for the participants in this study.

Randomised-controlled supplementation trials, where mothers were supplemented with PUFA during pregnancy, lactation or both, and the child's later cognitive performance was assessed, report varied conclusions^(38,39). Some studies have shown positive associations between LCPUFA and executive function, specifically in the domains of planning and working memory^(10,40). A study by Helland *et al.*⁽⁶⁾ found that children of mothers consuming LCPUFA supplements during pregnancy had higher intelligence quotient at 4 years of age than children whose mothers had not been supplemented. However, Ghys *et al.*⁽⁴¹⁾ found no association between cognitive performance at 4 years of age and phospholipid DHA and AA levels at birth. A recent meta-analysis has shown that PUFA supplementation associated positively with cognition only in PUFA-deficient participants⁽⁴²⁾. Additional differences among these studies include timing of supplementation, failure to measure blood levels of FA, a wide-array of cognitive outcome measures and a focus on n -3 FA rather than a full analysis of all FA.

It has been suggested that discrepant results from supplementation studies are due to genetic variation in the fatty acid desaturase (*FADS*) gene cluster^(43,44). Genetic differences in FA enzymes can alter how individuals process fats that they consume in their diet. When FA levels are measured in the blood, these enzymatic processes are accounted for automatically.



Therefore, measuring FA in blood is an improvement over measuring FA intake when it comes to determining the associations between specific fats and cognitive development. The direct analysis of blood FA avoids the potential confounding effect of genetic differences that may decrease blood levels of LCPUFA due to altered FA desaturase activity⁽⁴³⁾. Thus, supplementation studies that do not assess genetic variation or blood levels of FA miss this critical parameter.

Specificity of cognitive testing may also contribute to the discrepant results⁽³⁹⁾. For instance, a systematic review (that included individuals of all ages) reported that *n*-3 FA intake for more than 3 months did not affect executive function⁽⁴⁰⁾. In our study, current *n*-3 FA levels only tended to be associated with executive function. However, in a study by Colombo *et al.*⁽³¹⁾, LCPUFA supplementation in infancy improved executive function in later childhood, but the report did not include current FA levels. In the Colombo study, LCPUFA supplementation did not affect language and development at 18 months or spatial memory, simple inhibition or advanced problem solving at any age. Thus, it is evident that specific testing can lead to different conclusions.

This study has several strengths. Studies to date have focused on FA intake of mothers during pregnancy and lactation, the FA intake of babies through breast milk or formula or the FA intake of school-aged children. The current study assessed a set of twenty-five FA rather than focusing solely on the LCPUFA such as DHA and EPA. Some studies have measured erythrocyte or plasma FA in Tanzanian women and infants^(45–47). The current study addressed the gap between 4 and 6 years of age. This is an important time period to understand, because it is a time of major brain growth⁽³⁶⁾ and a time that the neurons of the prefrontal cortex continue to be myelinated from childhood into adolescence⁽¹³⁾. In addition, we report associations between current FA blood levels and the current cognitive performance of the children, providing a direct link between current FA status and cognition. Although Tanzanian children are routinely monitored for growth, there is no formal system to monitor cognitive development. Most studies that assess cognition in young children use the Bayley scales of development^(39,48–51), but our study utilised a specific test of executive function. Therefore, this study provides a first look at the associations between nutrition and executive function in Tanzanian children. The DCCS tasks are an excellent tool for assessing executive function in young children from a variety of backgrounds and experiences. This task can be customised to fit the cultural expectations of the population being analysed⁽¹²⁾. In this case, we customised the DCCS tasks to include animals and colours that were likely to be familiar to the children. In addition, the test was performed in the local language, Swahili. However, the paper-based format of testing was novel to the children because these children typically do not have access to paper before attending school at the age of 7 years. Consistent with previous investigations conducted in the USA⁽³⁰⁾ and Scotland⁽²⁹⁾, in our population >50% of those ≥48 months of age were able to successfully complete at least one DCCS task, regardless of their FA status. This suggests that our application of this modified DCCS in this population is valid. Further, another strength of the present investigation was the use of two common scoring approaches for the DCCS. For each scoring method, performance on the DCCS was similarly related to FA levels.

There are some limitations to our study. As this study was cross-sectional, all reported associations are correlative rather than causative. This study was conducted in one village in rural Tanzania, and thus the results are not generalisable to children residing in other areas of Tanzania or other areas of the world. The study was powered to detect effects of EFA. Therefore, the lack of association between cognitive function and individual long-chain *n*-3 FA such as DHA may have been due to the number of subjects analysed. During this study, blood samples were collected throughout the day. No fasting was required. This may increase variability in the whole-blood FA measurements. However, in this setting, differences are likely to be small because children from the village consume relatively similar and low-fat meals compared with children in other settings. In addition, we analysed whole-blood samples for lipids and were not able to differentiate among the source compartment of the lipid. Furthermore, blood levels of FA may not only indicate higher intake of that FA but also may reflect changes in the conversion of that FA to a longer-chain or desaturated FA. Children in this village had additional nutritional deficiencies, and we have corrected for those parameters for which we had data. Although socio-economic data were not collected from parents/caregivers, the population was relatively homogeneous in this regard⁽¹⁴⁾.

In summary, the results of this study suggest that whole-blood EFA levels are associated with cognition. Intervention trials that include assessment of whole-blood FA levels are required to determine the relationships between intake, blood levels and executive function in Tanzanian children.

Acknowledgements

The authors acknowledge the assistance of Mariana Ngowi and the village research teams who assisted during data collection.

This study was made possible by the generous support of American people through the United States Agency for International Development (USAID)-funded Innovative Agricultural Research Initiative project (iAGRI) (no. CA-621-A-00-11-00009-00). The opinions reported here are those of the authors and do not necessarily reflect the views of USAID or the United States Government.

T. J., S. S. C., J. K., M. B. P. and J. I. F. designed the study. T. J. and S. S. C. conducted the study. W. S. H., J. K., M. B. P. and J. I. F. provided essential materials necessary for the research. T. J., S. S. C., M. B. P. and J. I. F. analysed the data. All authors made contributions to the manuscript, but J. I. F. has the primary responsibility for the final content. All authors contributed to the critical interpretation and writing of the article and approved the final version.

No authors report conflicts of interest except W. S. H. whose laboratory (OmegaQuant Analytics, LLC) performed the DBS FA analyses.

Supplementary material

For supplementary material/s referred to in this article, please visit <http://dx.doi.org/doi:10.1017/S0007114516003494>

References

1. Carlson SJ, Fallon EM, Kalish BT, *et al.* (2013) The role of the omega-3 fatty acid DHA in the human life cycle. *JPEN J Parenter Enteral Nutr* **37**, 15–22.
2. Uauy R & Dangour AD (2006) Nutrition in brain development and aging: role of essential fatty acids. *Nutr Rev* **64**, 24–33.
3. Gomez-Pinilla F (2008) Brain foods: the effects of nutrients on brain function. *Nat Rev Neurosci* **9**, 568–578.
4. Briend A, Dewey KG & Reinhart GA (2011) Fatty acid status in early life in low-income countries – overview of the situation, policy and research priorities. *Matern Child Nutr* **7**, Suppl. 2, 141–148.
5. Lenroot RK, Gogtay N, Greenstein DK, *et al.* (2007) Sexual dimorphism of brain developmental trajectories during childhood and adolescence. *Neuroimage* **36**, 1065–1073.
6. Helland IB, Smith L, Saarem K, *et al.* (2003) Maternal supplementation with very-long-chain n-3 fatty acids during pregnancy and lactation augments children's IQ at 4 years of age. *Pediatrics* **111**, e39–e44.
7. Helland IB, Smith L, Blomén B, *et al.* (2008) Effect of supplementing pregnant and lactating mothers with n-3 very-long-chain fatty acids on children's IQ and body mass index at 7 years of age. *Pediatrics* **122**, 472–479.
8. Clandinin MTCJ & Van Aerde JEE (1989) Requirements of newborn infants for long chain polyunsaturated fatty acids. *Acta Paediatr Scand* **78**, 63–71.
9. Huffman SL, Harika RK, Eilander A, *et al.* (2011) Essential fats: how do they affect growth and development of infants and young children in developing countries? A literature review. *Matern Child Nutr* **7**, Suppl. 3, 44–65.
10. Sheppard KW & Cheatham CL (2013) Omega-6 to omega-3 fatty acid ratio and higher-order cognitive functions in 7- to 9-year-olds: a cross-sectional study. *Am J Clin Nutr* **98**, 659–667.
11. Cheatham CL, Colombo J & Carlson SE (2006) n-3 Fatty acids and cognitive and visual acuity development: methodologic and conceptual considerations. *Am J Clin Nutr* **83**, 1458S–1466S.
12. Zelazo PD (2006) The Dimensional Change Card Sort (DCCS): a method of assessing executive function in children. *Nat Protoc* **1**, 297–301.
13. Hughes D & Bryan J (2003) The assessment of cognitive performance in children: considerations for detecting nutritional influences. *Nutr Rev* **61**, 413–422.
14. Ntwenya JE, Kinabo J, Msuya J, *et al.* (2015) Dietary patterns and household food insecurity in rural populations of Kilosa district, Tanzania. *PLOS ONE* **10**, e0126038.
15. Ntwenya J, Kinabo J, Msuya J, *et al.* (2015) Household food insecurity and associated factors in rural communities: a case of Kilosa District, Tanzania. *African J Agricult Res* **10**, 4783–4794.
16. World Health Organization (2010) *WHO Anthro for Personal Computers, 3.2.2 2011 ed., pp. Software for Assessing Growth and Development of the World's Children (0 to 60 Months of Age)*. Geneva: WHO.
17. World Health Organization (2009) *WHO AnthroPlus for Personal Computers, pp. Software for Assessing Growth of the World's Children and Adolescents (5 to 19 years)*. Geneva: WHO.
18. Jumbe T, Comstock SS, Hahn SL, *et al.* (2016) Whole blood levels of the n-6 essential fatty acid linoleic acid are inversely associated with stunting in 2-to-6 year old Tanzanian Children: a cross-sectional study. *PLOS ONE* **11**, e0154715.
19. Harris WS, Varvel SA, Pottala JV, *et al.* (2013) Comparative effects of an acute dose of fish oil on omega-3 fatty acid levels in red blood cells versus plasma: implications for clinical utility. *J Clin Lipidol* **7**, 433–440.
20. Johnston DT, Deuster PA, Harris WS, *et al.* (2013) Red blood cell omega-3 fatty acid levels and neurocognitive performance in deployed U.S. Servicemembers. *Nutr Neurosci* **16**, 30–38.
21. Sarter B, Kelsey KS, Schwartz TA, *et al.* (2015) Blood docosahexaenoic acid and eicosapentaenoic acid in vegans: associations with age and gender and effects of an algal-derived omega-3 fatty acid supplement. *Clin Nutr* **34**, 212–218.
22. Marangoni F, Colombo C & Galli C (2004) A method for the direct evaluation of the fatty acid status in a drop of blood from a fingertip in humans: applicability to nutritional and epidemiological studies. *Anal Biochem* **326**, 267–272.
23. Bailey-Hall E, Nelson EB & Ryan AS (2008) Validation of a rapid measure of blood PUFA levels in humans. *Lipids* **43**, 181–186.
24. Siguel E (1998) Diagnosing essential fatty acid deficiency. *Circulation* **97**, 2580–2583.
25. Siguel EN, Chee KM, Gong JX, *et al.* (1987) Criteria for essential fatty acid deficiency in plasma as assessed by capillary column gas-liquid chromatography. *Clin Chem* **33**, 1869–1873.
26. Sauberlich HE (1999) *Laboratory Tests for the Assessment of Nutritional Status*, 2nd ed, Modern Nutrition. Boca Raton, FL: CRC Press LLC.
27. Zelazo PD, Muller U, Frye D, *et al.* (2003) The development of executive function in early childhood. *Monogr Soc Res Child Dev* **68**, vii–137.
28. Diamond A, Carlson SM & Beck DM (2005) Preschool children's performance in task switching on the dimensional change card sort task: separating the dimensions aids the ability to switch. *Dev Neuropsychol* **28**, 689–729.
29. Rennie DA, Bull R & Diamond A (2004) Executive functioning in preschoolers: reducing the inhibitory demands of the dimensional change card sort task. *Dev Neuropsychol* **26**, 423–443.
30. Kirkham NZ, Cruess L & Diamond A (2003) Helping children apply their knowledge to their behavior on a dimension-switching task. *Dev Sci* **6**, 449–476.
31. Colombo J, Carlson SE, Cheatham CL, *et al.* (2013) Long-term effects of LCPUFA supplementation on childhood cognitive outcomes. *Am J Clin Nutr* **98**, 403–412.
32. Cohen J (1988) *Statistical Power Analysis for the Behavioral Sciences*. Hillsdale, NJ: Lawrence Erlbaum Associates.
33. Bhargava A, Jukes M, Ngorosho D, *et al.* (2005) Modeling the effects of health status and the educational infrastructure on the cognitive development of Tanzanian schoolchildren. *Am J Hum Biol* **17**, 280–292.
34. Benjamini Y & Hochberg Y (1995) Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc, B* **57**, 289–300.
35. World Health Organization & United Nations International Childrens Emergency Fund (2003) *Global Strategy for Infant and Young Child Feeding*, no. resolution WHA55.25. Geneva: WHO.
36. Kuzawa CW, Chugani HT, Grossman LI, *et al.* (2014) Metabolic costs and evolutionary implications of human brain development. *Proc Natl Acad Sci U S A* **111**, 13010–13015.
37. Nakamura MT, Yudell BE & Loor JJ (2014) Regulation of energy metabolism by long-chain fatty acids. *Prog Lipid Res* **53**, 124–144.
38. Campoy C, Escolano-Margarit MV, Anjos T, *et al.* (2012) Omega 3 fatty acids on child growth, visual acuity and neurodevelopment. *Br J Nutr* **107**, Suppl. 2, S85–S106.
39. Sun H, Como PG, Downey LC, *et al.* (2015) Infant formula and neurocognitive outcomes: impact of study end-point selection. *J Perinatol* **35**, 867–874.



40. Jiao J, Li Q, Chu J, *et al.* (2014) Effect of *n*-3 PUFA supplementation on cognitive function throughout the life span from infancy to old age: a systematic review and meta-analysis of randomized controlled trials. *Am J Clin Nutr* **100**, 1422–1436.
41. Ghys A, Bakker E, Hornstra G, *et al.* (2002) Red blood cell and plasma phospholipid arachidonic and docosahexaenoic acid levels at birth and cognitive development at 4 years of age. *Early Hum Dev* **69**, 83–90.
42. Cooper RE, Tye C, Kuntsi J, *et al.* (2015) Omega-3 polyunsaturated fatty acid supplementation and cognition: a systematic review and meta-analysis. *J Psychopharmacol* **29**, 753–763.
43. Glaser C, Lattka E, Rzehak P, *et al.* (2011) Genetic variation in polyunsaturated fatty acid metabolism and its potential relevance for human development and health. *Matern Child Nutr* **7**, Suppl. 2, 27–40.
44. Lauritzen L, Fewtrell M & Agostoni C (2015) Dietary arachidonic acid in perinatal nutrition: a commentary. *Pediatr Res* **77**, 263–269.
45. Luxwolda MF, Kuipers RS, Koops JH, *et al.* (2014) Interrelationships between maternal DHA in erythrocytes, milk and adipose tissue. Is 1 wt% DHA the optimal human milk content? Data from four Tanzanian tribes differing in lifetime stable intakes of fish. *Br J Nutr* **111**, 854–866.
46. Kuipers RS, Luxwolda MF, Sango WS, *et al.* (2011) Postdelivery changes in maternal and infant erythrocyte fatty acids in 3 populations differing in fresh water fish intakes. *Prostaglandins Leukot Essent Fatty Acids* **85**, 387–397.
47. Luxwolda MF, Kuipers RS, Boersma ER, *et al.* (2014) DHA status is positively related to motor development in breastfed African and Dutch infants. *Nutr Neurosci* **17**, 97–103.
48. Sudfeld CR, McCoy DC, Fink G, *et al.* (2015) Malnutrition and its determinants are associated with suboptimal cognitive, communication, and motor development in Tanzanian Children. *J Nutr* **145**, 2705–2714.
49. Fahmida U, Htet MK, Adhiyanto C, *et al.* (2015) Genetic variants of FADS gene cluster, plasma LC-PUFA levels and the association with cognitive function of under-two-year-old Sasaknese Indonesian children. *Asia Pac J Clin Nutr* **24**, 323–328.
50. Qawasmi A, Landeros-Weisenberger A, Leckman JF, *et al.* (2012) Meta-analysis of long-chain polyunsaturated fatty acid supplementation of formula and infant cognition. *Pediatrics* **129**, 1141–1149.
51. Murray-Kolb LE, Rasmussen ZA, Scharf RJ, *et al.* (2014) The MAL-ED cohort study: methods and lessons learned when assessing early child development and caregiving mediators in infants and young children in 8 low- and middle-income countries. *Clin Infect Dis* **59**, Suppl. 4, S261–S272.