


RESEARCH ARTICLE

Effect of soybean and seaweed-based diets on growth performance, feed utilization, and gut microbiota of tilapia: A systematic review and meta-analysis

Leonildo dos Anjo Viagem^{1,2,3*}, Jean Nepomuscene Hakizimana¹, Cyrus Rumisha², Brunno da Silva Cerozi⁴, Gerald Misinzo^{1,5*}

1 SACIDS Africa Centre of Excellence for Infectious Diseases, SACIDS Foundation for One Health, Sokoine University of Agriculture, Morogoro, Tanzania, **2** Department of Animal, Aquaculture and Range Sciences, College of Agriculture, Sokoine University of Agriculture, Morogoro, Tanzania, **3** Department of Food and Agricultural Sciences, Rovuma University, Cabo Delgado, Mozambique, **4** Department of Animal Science, College of Agriculture, University of São Paulo, Piracicaba, São Paulo, Brazil, **5** Department of Veterinary Microbiology, Parasitology and Biotechnology, College of Veterinary Medicine and Biomedical Sciences, Sokoine University of Agriculture, Morogoro, Tanzania

* leonildo.viagem@sacids.org (LAV); gerald.misinzo@sacids.org (GM)



OPEN ACCESS

Citation: Viagem LdA, Hakizimana JN, Rumisha C, Cerozi BdS, Misinzo G (2024) Effect of soybean and seaweed-based diets on growth performance, feed utilization, and gut microbiota of tilapia: A systematic review and meta-analysis. PLoS ONE 19(7): e0293775. <https://doi.org/10.1371/journal.pone.0293775>

Editor: Amel Mohamed El Asely, Benha University, EGYPT

Received: October 18, 2023

Accepted: June 6, 2024

Published: July 24, 2024

Copyright: © 2024 Viagem et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its [Supporting Information](#) files.

Funding: This study was supported by the Partnership for Skills in Applied Sciences, Engineering, and Technology (PASET) through the Regional Scholarship and Innovation Fund (RSIF) awarded to LA Viagem to carry out Ph.D. studies at SACIDS Africa Centre of Excellence for Infectious Diseases, SACIDS Foundation for One Health,

Abstract

Tilapia, a significant aquaculture species globally, relies heavily on feed for its production. While numerous studies have investigated the impact of soybean and seaweed-based diets on tilapia, a comprehensive understanding remains elusive. This review aimed at evaluating and synthesizing the existing literature on these diets' effects, focusing on growth performance, feed utilization, and gut microbiota. A systematic search of databases was conducted using Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines and a total of 57 studies were included in the qualitative analysis and 24 in the meta-analysis. The results indicated that soybean-based diets, at a 59.4% inclusion level improved the Specific Growth Rate (SGR) of tilapia with an effect size of -2.14 (95% CI: -2.92, -1.37; $p < 0.00001$; $I^2 = 99\%$) and did not improve the feed conversion rate (FCR), as the effect size was 1.80 (95% CI: 0.72, 2.89; $p = 0.001$; $I^2 = 100\%$). For seaweed-based diets, at a 15.9% inclusion level did not improve SGR, with an effect size of -0.74 (95% CI: -1.70, 0.22; $p = 0.13$; $I^2 = 99\%$), and the FCR with an effect size of -0.70 (95% CI: -1.94, 0.54; $p = 0.27$; $I^2 = 100\%$). Regarding the gut microbiota, was noted a lack of studies meeting the inclusion criteria for tilapia. However, findings from studies on other farmed fishes suggested that soybean and seaweed-based diets could have diverse effects on gut microbiota composition and promote the growth of beneficial microbiota. This study suggests that incorporating soybean-based diets at 59.4% inclusion can improve the SGR of tilapia. Seaweed-based diets, while not demonstrating improvement in the analyzed parameters with an inclusion level of 15.9%, have the potential to contribute to the sustainability of the aquaculture industry when incorporated at lower levels.

Sokoine University of Agriculture, Morogoro, Tanzania. The funders had no role in the study design, collection, and analysis of the information, the decision to publish the final manuscript.

Competing interests: The authors have declared that no competing interests exist

Introduction

Tilapia are among the most important freshwater fish species worldwide and play a crucial role in global aquaculture production. As adaptable species to various environmental conditions, tilapia can be reared in a wide range of farming systems, including ponds, cages, and recirculation aquaculture systems [1–3]. The success of tilapia farming largely depends on the formulation of cost-effective and nutritionally balanced feed and an appropriate feeding management strategy that promotes optimal growth performance and feed utilization [4].

Traditional feed ingredients for tilapia include fishmeal and fish oil due to their high protein content balanced essential amino acids profile and omega-3 [5–8], but they are expensive and can contribute to the overfishing of wild fish stocks [9–11]. Therefore, there is a growing interest in exploring alternative protein sources for fish feed, such as soybean and seaweed. Soybean is a cost-effective, and widely available plant protein source [12, 13], while seaweeds offer unique nutritional properties and are natural forage for fish [14]. Both have been investigated as partial or total replacements to improve the growth performance and feed efficiency of tilapia. Their inclusion in this study can provide a comprehensive understanding of their individual nutritional properties and highlight their potential as feed options for tilapia. However, there are challenges in including seaweed and soybeans in tilapia diets, such as mismatches in fatty acids and amino acids profile, leading to sub-optimal growth and health [15, 16]. Additionally, soybean and seaweed contain antinutritional factors such as saponins, tannins, phytic acid, lectins, protease inhibitors, and amylase inhibitors [16–18], which may negatively affect nutrient digestibility, palatability, and overall feed performance.

Apart from the nutritional aspects, the gut microbiota plays a crucial role in the overall health and growth of fish [19–21] and can participate in modulating feeding behavior and efficiency [22, 23]. Dietary interventions can influence the composition and activity of the gut microbiota, thereby influencing the host's metabolism, immune system, and overall performance [24–27].

While individual studies have investigated the effects of soybean and seaweed-based diets on tilapia, a comprehensive understanding of their impact on growth performance, feed utilization, and gut microbiota is lacking. A systematic review and meta-analysis are essential to synthesize the available evidence and provide a comprehensive overview of the effects of these alternative feed ingredients on tilapia. Therefore, the present study evaluated and synthesized the existing literature on the effects of soybean and seaweed-based diets on growth performance, feed utilization, and gut microbiota of tilapia to achieve sustainability in aquaculture practices.

Methods

Study design

This study was conducted according to Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines (S1 Table). The protocol related to this systematic review was registered on Open Science Framework (osf.io/zg72m) with registration DOI: <https://doi.org/10.17605/OSF.IO/ZG72M>.

Data source and search strategy

A systematic literature search was conducted using online databases including Google Scholar, PubMed, Wiley Online Library, and ScienceDirect, to identify relevant studies published between 2000 and 2022. The search terms were combined as follows: “seaweed” OR “macroalgae” NOT “microalgae” AND “growth performance” AND “feed utilization” AND “tilapia”; “soybean” NOT “soybean fermented” AND “growth performance” AND “feed utilization” AND “tilapia”; “seaweed” OR “macroalgae” NOT “microalgae” AND “gut microbiota” OR

“intestinal microbiota” AND “tilapia” OR “other fish” NOT “human”; “soybean” NOT “soybean fermented” AND “gut microbiota” OR “intestinal microbiota” AND “tilapia” OR “other fish” NOT “human”. Additionally, a manual search of references cited in relevant articles was performed to identify additional studies that may have been missed in the database search. The search strategies and the respective number of studies from the four databases are detailed in the [S2 Table](#).

Inclusion and exclusion criteria

In this systematic review, studies published from January 2000 to December 2022 were included based on the following criteria: (1) articles written in English; (2) articles focusing on tilapia production; (3) studies utilizing various soybean-based diets such as soybean defatted, soybean meal, full-fat soybean, soybean milk, and soybean protein; (4) studies utilizing seaweed-based diets, including seaweed extract and various species of green, brown, red seaweed ([S3 Table](#)) to partially or totally replace fish meal/oil; (5) full-text available; (6) studies published in peer-reviewed journals; (7) studies reporting SGR and FCR with their respective standard error or standard deviation, and (8) articles examining the effect of soybean and seaweed-based diets on intestinal microbiota of other fish than tilapia. Excluded from consideration were: (1) studies not written in English; (2) studies not related to the use of soybean and seaweed as tilapia feed; (3) articles not published in peer-reviewed scientific journals; (4) articles not available in full-text format; (5) studies that used microorganisms to ferment soybean/seaweed; and (6) studies utilizing prebiotics in tilapia/fish production that were not derived from soybean or seaweed; (7) article that did not describe the composition of the gut microbiota; (8) studies lacking fish meal or fish oil as control group.

Selection process

The screening of all papers was done in the Rayyan tool for systematic literature review (<https://www.rayyan.ai/>). After downloading the article in the databases (in RIS format), the files were imported to Rayyan to check for duplicates, screening the title, abstract, and full text by two independent reviewers (Leonildo dos Anjo Viagem-LAV and Jean Nepomuscen Hakimimana-JNH).

Data extraction

Data were extracted from each included study using a standardized data extraction form created in Microsoft Excel 2013 (Microsoft Corporation, Washington, USA). This extraction process was carried out independently by two reviewers (LAV and JNH). The extracted data encompassed basic information such as author and publication year, experimental details including sample size and experiment duration, the inclusion level of the diets (both applied and recommended), and information about the different experimental groups, namely the soybean-based diet group, seaweed-based diet group, and control group (fish meal or fish oil). Parameters extracted for these groups included specific growth rate (SGR), and feed conversion ratio (FCR), along with their corresponding standard mean or standard error. Additionally, details regarding the types of soybean and seaweed species utilized in the experiments were recorded ([S3 Table](#)).

Risk of bias in included studies

The study quality and risk of bias were evaluated using the Cochrane risk-of-bias tool in Cochrane Collaboration’s software Review Manager Version 5.4.1 (RevMan 5.4.1). This tool

contains seven domains: (1) random sequence bias (selection bias), (2) allocation concealment (selection bias), (3) blinding of participants and personnel (performance bias), (4) blinding of outcome assessment (detection bias), (5) incomplete outcome data (attrition bias), (6) selective reporting (reporting bias), and (7) other biases [28]. This systematic review focused on five domains, as the blinding of participants and personnel (performance bias) and the blinding of outcome assessment (detection bias) are not applicable to aquaculture studies. In the domain of other bias was analyzed the information related to sample size and experiment duration.

Meta-analysis

A meta-analysis was conducted to analyze the effect of soybean and seaweed on the SGR and FCR of tilapia. Effect sizes were calculated, and a random-effects model was employed. Publication bias was assessed using funnel plots of the effect of soybean on SGR and FCR (S1 Fig) and seaweed (S2 Fig). The heterogeneity of the included studies was assessed by the I-squared statistic test and was classified as low (<25%); moderate (25–50%) and high (>50%) [29]. The subgroup analysis and sensitivity were performed to explore possible causes of heterogeneity in the study. All statistical analyses were conducted in RevMan 5.4.1 and the variables were analyzed as continuous via standard mean difference with a 95% confidence interval (CI).

Results

Summary of included studies

Initially, a total of 17,991 studies were collected from the databases and 8 from other sources (Fig 1). After removing duplicates, 12,548 remained for screening. Based on eligibility criteria, 11,369 studies were excluded after titles and abstracts screening. The majority of the excluded studies focused on other animals than fish ($n = 5,805$) followed by those not addressing the topic of tilapia ($n = 2,853$) and seaweed or soybean ($n = 2,351$). Then, 1,179 full texts were assessed and 1,122 were excluded because most studies used probiotics to ferment soybean/seaweed ($n = 910$) and due to a lack of data on Specific Growth Rate or Feed Conversion Rate ($n = 122$). After full-text screening, 57 studies were included in the qualitative synthesis and 24 were included in the meta-analysis.

The included studies described that the fish were in good health and were purchased from companies that regularly conduct health inspections. These studies exhibited variations in experimental design, duration of the experiment, manner of processing soybeans, species of seaweed used in the experiment, and dietary inclusion level. In all the included papers, soybeans and seaweed were studied individually and in this review were analyzed separately.

Quality and risk of bias assessment

The quality and risk of bias of included studies were assessed through the Cochrane criteria showing positive results for allocation concealment (selection bias), incomplete outcome data (attrition bias), selective reporting (reporting bias), and other bias (sample size and experiment duration) (Fig 2A and 2B). The evaluation of the studies revealed unclear risk bias for domain random sequence bias (selection bias) because most of the studies reported the random method but the sequence generation was not described. The studies Abdel-Warith et al., (2013), El-Saidy et al., (2010) and Sharda et al., (2017) (Fig 3A) and Stadtlander et al., (2012), Radwan et al., (2022) and Eissa et al., (2021) (Fig 3B) revealed high-risk bias for random sequence bias (selection bias). The study by Stadtlander et al., (2012) showed unclear bias for other bias because the sample size was not enough for this systematic review but the experiment duration was reported.




Fig 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flowchart of the study selection process. *n* represents the number of studies.

<https://doi.org/10.1371/journal.pone.0293775.g001>

Effect of soybean on growth performance and feed utilization

In this study, Specific Growth Rate (SGR) and Feed Conversion Rate (FCR) were used as parameters of growth performance and feed utilization, respectively. Among the studies




Fig 2. The risk of bias graph shows the review author's judgment about each risk of bias item presented as percentages across all included studies on soybean (A) and seaweed (B). The quality and risk of bias was assessed using the Cochrane risk-of-bias tool in Cochrane Collaboration's software RevMan 5.4.1.

<https://doi.org/10.1371/journal.pone.0293775.g002>




Fig 3. Risk of bias summary showing review author's judgment about each risk of bias item for each included study on soybean (A) and seaweed (B). The quality and risk of bias was assessed using the Cochrane risk-of-bias tool in Cochrane Collaboration's software RevMan 5.4.1.

<https://doi.org/10.1371/journal.pone.0293775.g003>

included in the meta-analysis, twelve [11, 16, 30–39] described the effects of soybean on growth performance and feed utilization of tilapia (*Oreochromis niloticus* and *Sarotherodon melanotheron*) and involved 8,055 fish and 53 treatments. One article (Koumi et al., 2011) studied 2 species of tilapia, and the analysis was performed separately. The results showed that the random pooled effect size of SGR was -2.14 (95% CI: $-2.92, -1.37$; $p < 0.00001$; $I^2 = 99\%$) (Fig 4A) and for FCR was 1.80 (95% CI: $0.72, 2.89$; $p = 0.001$; $I^2 = 100\%$) (Fig 4B). The average inclusion level of soybeans in the diets, calculated from the recommendation in each analyzed study (S3 Table), was found to be 59.4%. These findings suggest that including soybeans at this level can enhance the SGR but does not lead to an improvement in FCR.

Effect of seaweed on growth performance and feed utilization

Among the studies included in the meta-analysis, twelve [8, 40–50] described the effects of seaweed on growth performance and feed utilization of *O. niloticus* and hybrid red tilapia (*O. niloticus* and *Oreochromis mossambicus*), involving 5,657 fish and 55 treatments of different species including rede seaweed (*Laurencia obtusa*, *Pterocladia capillacea*, *Gracilaria arcuata*, *Porphyra yezoensis*, *Gracilaria vermiculophylla*, and *Porphyra dioica*); green seaweed (*Ulva lactuca*, *Enteromorpha flexuosa*, and *Ulva fasciata*) and brown seaweed (*Taonia atomaria* and *Cystoseira myrica*) (S3 Table). Two papers (Silva et al., 2014 and Eissa et al., 2020) each studied three species of seaweed while one paper (Khalafalla and El-Hais, 2015) studied 2 species of seaweed, and the analysis was performed separately for each species. The results showed that the random pooled effect size of SGR was -0.74 (95% CI: $-1.70, 0.22$; $p = 0.13$; $I^2 = 99\%$) (Fig 5A), and for FCR was -0.70 (95% CI: $-1.94, 0.54$; $p = 0.27$; $I^2 = 100\%$) (Fig 5B). The inclusion




Fig 4. Forest plot showing the effect of soybean on specific growth rate (A) and feed conversion rate (B) of tilapia. The analysis was performed on RevMan 5.4.1, based on twelve studies and the effect size index is the standardized mean difference. The random-effects model was employed for this analysis and the studies in the analysis are assumed to be a random sample from a universe of potential studies. The variables were analyzed as continuous via standard mean difference with a 95% CI. **O.n:** *Oreochromis niloticus*; **S.m:** *Sarotherodon melanotheron*; **CI:** Confidence Interval; **Tau²:** tau-squared; **df:** difference; **Chi²:** chi-squared; **P:** p-value; **I²:** squared statistic test; **Z:** z-value; **SD** and **Std:** Standard Deviation; *same study but carried out on two species of tilapia (*O. niloticus* and *S. melanotheron*).

<https://doi.org/10.1371/journal.pone.0293775.g004>

level of seaweed was 15,9%. These results indicate that the inclusion level of seaweed at 15,9% did not improve the SGR and FCR of tilapia.

Effect of soybean and seaweed on the gut microbiota of tilapia

The search conducted in the databases using the search strategies defined in this systematic review on tilapia was not able to identify studies that met the inclusion criteria on this specific topic. Most of the studies were done on other farmed fish (Table 1), showing that soybean and seaweed could have different effects on gut microbiota composition and diversity.

Discussion

Effect of soybean and seaweed-based diet on SGR and FCR

The findings of this review suggest an enhancement in SGR among tilapia fed with soybean-based diets. This improvement could be attributed to soybean's high protein content and relatively balanced amino acid profile [33, 34]. The findings also reveal that soybean did not improve FCR, likely due to the presence of antinutritional factors such as protease inhibitors, lectins, phytic acid, saponins, phytoestrogens, antivitamin, and allergens [16, 17]. These factors may compromise the palatability of soybeans and thus affect their effectiveness in improving FCR. The results from studies examining the impact of soybeans suggest that growth performance and feed utilization of tilapia tend to decline as the inclusion levels of soybeans




Fig 5. Forest plot showing the effect of seaweed on specific growth rate (A) and feed conversion rate (B) of tilapia. The analysis was performed on RevMan 5.4.1, based on seventeen studies and the effect size index is the standardized mean difference. The random-effects model was employed for this analysis and the studies in the analysis are assumed to be a random sample from a universe of potential studies. The variables were analyzed as continuous via standard mean difference with a 95% CI. **GRA:** *Gracillaria*; **UV:** *Ulva*; **LA:** *Laurencia*; **POR:** *Porphyra*; **CY:** *Cystoseira*; **CI:** Confidence Interval; **Tau²:** tau-squared; **df:** difference; **Chi²:** chi-squared; **P:** p-value; **I²:** squared statistic test; **Z:** z-value; **SD** and **Std:** Standard Deviation; ** same study but carried out on different species of seaweed.

<https://doi.org/10.1371/journal.pone.0293775.g005>

increase, the ideal level of inclusion being up to 50% [31, 35]. Similar studies conducted on other fish species such as *Liza haematocheila* [67]; *Heteropneustes fossilis* [68]; *Takifugu obscurus* [69]; *Clarias gariepinus* [70]; *Solea aegyptiaca* [71]; *Sparus aurata* [72]; also indicate a decrease in growth parameters with higher levels of soybean inclusion level. However, determining the optimal inclusion level and type of soybean (e.g., defatted, fermented, full fat) requires further investigation since the studies included in the meta-analysis utilized different inclusion levels and types of soybeans. On the other hand, the results do not support the use of seaweed as an alternative protein source for tilapia, as it did not improve the SGR and FCR. However, it should be noted that the studies included in the meta-analysis used different species of seaweeds and different levels of inclusion.

Several studies have demonstrated that seaweed inclusion levels of up to 20% can effectively replace fishmeal in tilapia diets without compromising the growth performance and feed utilization [40, 41, 43, 48, 50, 73]. Similar outcomes have been observed in other fish species as well [74–76]. The bioactive compounds present in soybeans and seaweed, such as Non-Starch Polysaccharides (NSPs) and Antinutritional Factors (ANFs) like saponins, lectins, and phytates, play significant roles in influencing fish digestibility and metabolic functions [77, 78]. Excessive NSPs may cause gut inflammation and reduce energy availability, while ANFs can interfere with digestive enzymes and nutrient absorption [79]. However, proper processing methods such as heat and enzymatic treatment, bioprocessing, and fermentation of these diets can mitigate these effects [18, 79, 80]. Furthermore, phytochemicals like isoflavones in soybeans possess antioxidant properties, although their impact may vary among fish species [81].

Table 1. The most abundant gut microbiota in other fish than tilapia fed with soybean and seaweed-based diet.

Type of diet	Inclusion level	Fish species	Composition of gut microbiota	Reference
Soybean meal (SBM)	25%	Atlantic salmon (<i>Salmo salar</i>)	<i>Vibrio</i> spp., <i>Arthrobacter agillis</i> , <i>Brachybacterium</i> spp., <i>Kocuria carniphila</i> and <i>Rhodococcus</i> spp	[51]
Green Seaweed (<i>Ulva ohnoi</i>)	5%	Flatfish (<i>Solea senegalensis</i>)	<i>Vibrionaceae</i> and <i>Shewanella</i>	[52]
Red Seaweed (<i>Gracilaria gracilis</i>)	8%	European seabass (<i>Dicentrarchus labrax</i>)	<i>Acinetobacter</i> , <i>Corynebacterium</i> , <i>Ilumatobacter</i> , <i>Kordia</i> , <i>Methylobacterium</i> , <i>Polaribacter</i> , <i>Pseudomonas</i> and <i>Sulfitobacter</i>	[53]
Red seaweed (<i>Asparagopsis taxiformis</i>)	3%	Atlantic salmon (<i>Salmo salar</i>)	<i>Aeromodaceae</i>	[54]
Soybean meal	10%	Atlantic salmon (<i>Salmo salar</i>)	<i>Psychrobacter okhotskensis</i> , <i>Arthrobacter rhombi</i> , <i>Arthrobacter psychrolactophilus</i> , <i>Psychrobacter cibarius</i> and <i>Psychrobacter fozii</i>	[55]
Red Seaweed (<i>Gracilaria cornea</i>) and green seaweed (<i>Ulva rigida</i>)	5%, 15% and 25%	Gilthead seabream (<i>Sparus aurata</i>) juveniles	Gracillaria (15% and 25%): <i>Pseudomonas</i> and <i>Lactobacillus</i> ; Ulva (5% and 15%): <i>Lactobacillus</i> and <i>Sphingomonas</i> ; Ulva (25%): <i>Comamonas aquatica</i>	[56]
Soybean meal (SBM)	20%	Atlantic salmon (<i>Salmo salar</i> L.)	<i>Lactococcus lactis</i> subsp. <i>lactis</i> , <i>Weissella confusa</i> , and <i>Photobacterium phosphoreum</i>	[57]
Soybean meal (SBM)	50%	Atlantic salmon (<i>Salmo salar</i>)	<i>Aeromonas</i> and <i>Sporosarcina aquimarina</i>	[58]
Soybean meal (SBM)	50%	Rainbow trout (<i>Oncorhynchus mykiss</i>)	<i>Psychrobacter</i> spp and <i>Saccharomyces</i> spp	[59]
Soybean Meal (SBM)	20%– 100%	<i>Carassius auratus gibelio</i> ♀ + <i>Cyprinus carpio</i> ♂	<i>Cetobacterium</i> (10.53–17.46%) and <i>Aeromonas</i> (4.68–13.88%)	[60]
Green seaweed (<i>Ulva rigida</i>)	10%	Atlantic cod (<i>Gadus morhua</i>)	<i>Ruminococcaceae</i> spp., <i>Shewanella</i> , <i>Tyzerella</i> , <i>Rikenellaceae</i> spp., <i>Peptococcaceae</i> , <i>Bacteroides</i> and <i>Fusobacterium</i> .	[61]
Green seaweed (<i>Ulva rigida</i>)	25%	<i>Sparus aurata</i> juveniles	<i>Lactobacillus delbrueckii</i>	[62]
Soybean meal (SBM)	45%	Juvenile rainbow trout (<i>Oncorhynchus mykiss</i>)	<i>Aeromonas</i> , <i>Sphingomonas</i> and <i>Chryseomonas</i>	[63]
Red seaweed (<i>Gracilaria gracilis</i>)	2.5% and 5%	European seabass (<i>Dicentrarchus labrax</i>)	<i>Sphingomonas</i> , <i>Photobacterium</i> , <i>Staphylococcus</i> and <i>Vibrio</i>	[64]
Soybean Protein Isolate (SPI)	25% and 50%	Spotted seabass (<i>Lateolabrax maculatus</i>)	<i>Prevotella</i> , <i>Succiniclaticum</i> , and <i>Desulfovibrio</i>	[65]
Green seaweed (<i>Ulva ohnoi</i>)	5%	<i>Solea senegalensis</i>	<i>Pseudomonas</i> and <i>Mycoplasmataceae</i>	[66]

<https://doi.org/10.1371/journal.pone.0293775.t001>

Additionally, certain seaweed compounds, like polysaccharides may promote gut health, enhancing overall digestion and nutrient absorption [82, 83]. However, the specific impact can vary depending on the type and amount of seaweed, as well as the fish species involved [15, 84]. For instance, several studies have demonstrated that the green seaweed *Ulva* spp, when included at levels of up to 5%, can enhance the growth performance of certain carnivorous fish [85, 86], and omnivorous fish like tilapia of up to 20% of inclusion [87]. In addition to soybeans and seaweed, plant-based diets such as sunflower, canola, cottonseed, Moringa, and *Azolla* meals have shown promise in feeding tilapia and other fish [88–92]. *Azolla*, noted for its exceptional amino acid profile, exhibits the highest nutritional index in tilapia feed [93]. In the data analysis, heterogeneity was observed among the included studies, potentially due to variations in experimental design, number of fish used in each study, type of soybean, and species of seaweed used across different studies. Subgroup analyses were performed to explore potential sources of heterogeneity, but the findings were inconclusive.

Effect of soybean and seaweed-based diet on gut microbiota

In this review, it was not possible to find studies that describe the effect of soybean or seaweed on the gut microbiota of tilapia. However, studies in other fish species have shown diverse effects on the gut microbiota composition. For instance, beneficial bacteria like *Shewanella*

were observed in Flatfish and Atlantic cod fed with *Ulva* spp [52, 61], while *Lactobacillus* spp were found in Gilthead seabream fed with *Gracilaria cornea* and *Ulva rigida* [56, 62] along with *Lactococcus* spp [57]. Additionally, *Psychrobacter* spp, and *Arthrobacter* spp were noted in fish fed with soybean meal [51, 55, 59]. These microorganisms are known to possess probiotic properties, which could potentially enhance nutrient absorption and disease resistance [94–97]. The seaweed and soybean dietary components contain bioactive compounds such as isoflavones, saponins, and oligosaccharides in soybean [98–100], and polysaccharides, polyphenols, phycobiliproteins, and phlorotannins in seaweed [101–103]. These compounds can exert prebiotic effects, influencing the composition of the gut microbiota [104]. Prebiotics are essentially non-digestible substances that, when consumed, foster beneficial changes in the gut microbiome, promoting the development of a more favorable microbial community [105]. This shift can have positive implications for various aspects of health, including digestive function, immune system modulation, and antimicrobial properties, inhibiting the growth of potentially harmful bacteria [106–108]. It is important to note that the specific effects of soybean and seaweed on fish gut microbiota can vary depending on various factors, such as the fish species, the inclusion levels of these ingredients, and the species of seaweed. Studies investigating the utilization of various woody forages such as *Moringa oleifera*, fermented *M. oleifera*, *Folium mori*, and fermented *F. mori* meals in tilapia farming have indicated significant variations in gut microbiota composition [109, 110]. These findings suggest that diets incorporating soybean, seaweed, and other plant-based diets may have diverse effects on the diversity and composition of fish intestinal microbiota.

Implications for tilapia production

The inclusion of soybean and seaweed-based diets into tilapia farming can contribute to the sustainability of the aquaculture industry by reducing reliance on fishmeal and fish oil. The improved SGR observed in tilapia fed with a soybean diet indicates the potential for cost-effectiveness and viability in tilapia production. Regarding seaweed-based diets, although they may not directly improve growth performance and feed utilization parameters in tilapia, they offer alternative feed options due to their rich content of bioactive compounds. These compounds can have beneficial effects on fish growth and health. Additionally, the modulation of gut microbiota by soybean and seaweed-based diets can improve tilapia health and nutrient utilization leading to more sustainable and cost-effective tilapia production.

Limitations and future research directions

Despite the valuable insights gained from this study, limitations should be considered: (1) the included studies may have exhibited variations in experimental design, type of soybean and seaweed used, leading to heterogeneity in the data, (2) limitations for some outcomes, particularly effect of soybean and seaweed on gut microbiota of tilapia. For future research, it is suggested: (1) investigations into the optimal inclusion levels of soybean and seaweed-based ingredients and exploring the potential seaweed species for tilapia diets should be conducted to maximize growth performance and feed utilization; (2) long-term studies are needed to evaluate the potential impact of these diets on the gut microbiota and its implications for growth performance, nutrient utilization, and the health of tilapia and (3) investigate the effect of those diets in other species of tilapia.

Conclusion

This systematic review and meta-analysis provide valuable insights into the effects of soybean and seaweed-based diets on tilapia production. The results of the meta-analysis indicated that

the soybean inclusion level of 59.4% improved SGR but did not improve the FCR of tilapia. On the other hand, the seaweed-based diet did improve both SGR and FCR at an inclusion level of 15.9%. Based on the search strategies defined in this review, it was not possible to find studies that met the inclusion criteria for the gut microbiota of tilapia. However, studies on soybean and seaweed in other farmed fish have shown diverse effects on the gut microbiota composition and have the potential to enhance the growth of beneficial gut microbiota and improve fish health. This information is valuable for tilapia production as it offers an alternative dietary strategy for optimizing their growth.

Supporting information

S1 Table. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 checklist.

(DOCX)

S2 Table. Complete database search strategies and the number of articles accessed in each database.

(DOCX)

S3 Table. Data extraction from eligible studies on the effect of soybean and seaweed-based diets on tilapia production.

(XLSX)

S1 Fig. Publication bias represented by funnel plot of individual studies about the effect of soybean on SGR (A) and FCR (B).

(TIF)

S2 Fig. Publication bias represented by funnel plot of individual studies about the effect of seaweed on SGR (A) and FCR (B).

(TIF)

Author Contributions

Conceptualization: Leonildo dos Anjo Viagem.

Data curation: Leonildo dos Anjo Viagem, Jean Nepomuscene Hakizimana.

Formal analysis: Leonildo dos Anjo Viagem.

Methodology: Leonildo dos Anjo Viagem.

Supervision: Cyrus Rumisha, Brunno da Silva Cerozi, Gerald Misinzo.

Writing – original draft: Leonildo dos Anjo Viagem.

Writing – review & editing: Jean Nepomuscene Hakizimana, Cyrus Rumisha, Brunno da Silva Cerozi, Gerald Misinzo.

References

1. Arumugam M, Jayaraman S, Sridhar A, Venkatasamy V, Brown PB, Abdul Kari Z, et al. Recent Advances in Tilapia Production for Sustainable Developments in Indian Aquaculture and Its Economic Benefits. *Fishes*. 2023; 8(4):1–31.
2. FAO. Social and economic performance of tilapia farming in Brazil. Junning Cai KQ and NH, editor. Vol. 1130, Social and economic performance of tilapia farming in Brazil. Rome, Italy: FAO; 2017. 144 p.
3. Prabu E, Rajagopalsamy CBT, Ahilan B, Jeevagan IJMA, Renuhadevi M. Tilapia—An Excellent Candidate Species for World Aquaculture: A Review. *Annu Res Rev Biol*. 2019; 31(3):1–14.

4. Ng WK, Romano N. A review of the nutrition and feeding management of farmed tilapia throughout the culture cycle. *Rev Aquac.* 2013; 5(4):220–254.
5. El-Sayed AFM. Nutrition and feeding. *Tilapia Culture.* 2020. 135–172 p.
6. Norambuena F, Hermon K, Skrzypczyk V, Emery JA, Sharon Y, Beard A, et al. Algae in fish feed: Performances and fatty acid metabolism in juvenile Atlantic Salmon. *PLoS One.* 2015; 10(4):1–17. <https://doi.org/10.1371/journal.pone.0124042> PMID: 25875839
7. Yildirim-Aksoy M, Lim C, Li MH, Klesius PH. Interaction between dietary levels of vitamins C and E on growth and immune responses in channel catfish, *Ictalurus punctatus* (Rafinesque). *Aquac Res.* 2008 Aug 1; 39(11):1198–1209.
8. Younis ESM, Al-Quffail AS, Al-Asgah NA, Abdel-Warith AWA, Al-Hafedh YS. Effect of dietary fish meal replacement by red algae, *Gracilaria arcuata*, on growth performance and body composition of Nile tilapia *Oreochromis niloticus*. *Saudi J Biol Sci.* 2018; 25(2):198–203.
9. Naylor RL, Hardy RW, Bureau DP, Chiu A, Elliott M, Farrell AP, et al. Feeding aquaculture in an era of finite resources. *Proc Natl Acad Sci U S A.* 2009; 106(36):15103–15110. <https://doi.org/10.1073/pnas.0905235106> PMID: 19805247
10. Olsen RL, Hasan MR. A limited supply of fishmeal: Impact on future increases in global aquaculture production. *Trends Food Sci Technol.* 2012; 27(2):120–128.
11. Zhao H, Jiang R, Xue M, Xie S, Wu X, Guo L. Fishmeal can be completely replaced by soy protein concentrate by increasing feeding frequency in Nile tilapia (*Oreochromis niloticus* GIFT strain) less than 2 g. *Aquac Nutr.* 2010; 16(6):648–653.
12. Kook MC, Cho SC, Hong YH, Park H. *Bacillus subtilis* fermentation for enhancement of feed nutritive value of soybean meal. *J Appl Biol Chem.* 2014; 57(2):183–188.
13. Yun A, Kim J, Jeong HS, Lee KW, Kim HS, Kim PY, et al. Inclusion effect of soybean meal, fermented soybean meal, and *Saccharina japonica* in extruded pellet for juvenile abalone (*Haliotis discus*, Reeve 1846). *Fish Aquat Sci.* 2018; 21(1):1–8.
14. Rajauria G. Seaweeds: a sustainable feed source for livestock and aquaculture. In: Tiwari B, TroyD (eds) *Seaweed sustainability: food and non-food applications.* Seaweed Sustainability. Elsevier Inc.; 2015. 389–420 p.
15. Pereira R, Valente LMP, Sousa-Pinto I, Rema P. Apparent nutrient digestibility of seaweeds by rainbow trout (*Oncorhynchus mykiss*) and Nile tilapia (*Oreochromis niloticus*). *Algal Res.* 2012; 1(1):77–82.
16. Abdel-Warith A-WA-m., Younis EM, Al-Asgah NA. Influence of Dietary Inclusion of Full-Fat Soybean Meal and Amino Acids Supplementation on Growth and Digestive Enzymes Activity of Nile Tilapia, *Oreochromis niloticus*. *Turkish J Fish Aquat Sci.* 2013;13:69–77.
17. Hussain SM, Afzal M, Rana SA, Javid A, Iqbal M. Effect of phytase supplementation on growth performance and nutrient digestibility of *Labeo rohita* fingerlings fed on corn gluten meal-based diets. *Int J Agric Biol.* 2011; 13(6):916–922.
18. Samtiya M, Aluko RE, Dhewa T. Plant food anti-nutritional factors and their reduction strategies: an overview. *Food Prod Process Nutr.* 2020; 2(1):1–14.
19. Huang Q, Sham RC, Deng Y, Mao Y, Wang C, Zhang T, et al. Diversity of gut microbiomes in marine fishes is shaped by host-related factors. *Mol Ecol.* 2020; 29(24):5019–5034. <https://doi.org/10.1111/mec.15699> PMID: 33084100
20. Nie L, Zhou QJ, Qiao Y, Chen J. Interplay between the gut microbiota and immune responses of ayu (*Plecoglossus altivelis*) during *Vibrio anguillarum* infection. *Fish Shellfish Immunol.* 2017; 68:479–487.
21. Xiong JB, Nie L, Chen J. Current understanding of the roles of gut microbiota in fish disease and immunity. *Zool Res.* 2019; 40(2):70–76.
22. Leitão-Gonçalves R, Carvalho-Santos Z, Francisco AP, Fioreze GT, Anjos M, Baltazar C, et al. Commensal bacteria and essential amino acids control food choice behavior and reproduction. *PLoS Biol.* 2017; 15(4):1–29. <https://doi.org/10.1371/journal.pbio.2000862> PMID: 28441450
23. Yamazaki Y, Meirelles PM, Mino S., Suda KW., Hattori M, Thompson FL., et al. Individual *Apostichopus japonicus* fecal microbiome reveals a link with polyhydroxybutyrate producers in host growth gaps. *Sci Rep.* 2016; 6(February):1–10
24. Huang C, Liao W. Seasonal Variation in Gut Microbiota Related to Diet in *Fejervarya limnocharis*. *Animals.* 2021;1–15.
25. Piazzon MC, Caldach-Giner JA, Fouz B, Estensoro I, Simó-Mirabet P, Puyalto M, et al. Under control: how a dietary additive can restore the gut microbiome and proteomic profile, and improve disease resilience in a marine teleostean fish fed vegetable diets. *Microbiome.* 2017; 5(1):1–23.
26. Ringø E, Zhou Z, Vecino JLG, Wadsworth S, Romero J, Kroghdahl Å, et al. Effect of dietary components on the gut microbiota of aquatic animals. A never-ending story? *Aquac Nutr.* 2016 Apr 1; 22(2):219–282.

27. Wang T, Wu HX, Li WJ, Xu R, Qiao F, Du ZY, et al. Effects of dietary mannan oligosaccharides (MOS) supplementation on metabolism, inflammatory response, and gut microbiota of juvenile Nile tilapia (*Oreochromis niloticus*) fed with high carbohydrate diet. *Fish Shellfish Immunol*. 2022 Nov; 130:550–559.
28. Higgins J, Green S (Sally E, Cochrane Collaboration. Assessing risk of bias in included studies Cochrane handbook for systematic reviews of interventions. Cochrane, editor. 2017. 2017. 1p.
29. Julian P T Higgins, Simon G Thompson, Jonathan J Deeks DGA. Measuring inconsistency in meta-analyses. *BMJ* 2003;327. 2003;327(2):557–560.
30. Agbo NW, Madalla N, Jauncey K. Mixtures of oilseed meals as dietary protein sources in diets of juvenile Nile tilapia (*Oreochromis niloticus* L.). *J Sci Technol*. 2016; 35(3):11–24.
31. Ahmad N, Siddiqui PJA, Khan KM, Ali A, Tahir M, Akbar NU, et al. Effects of partial substitution of fish-meal by soybean meal in Nile tilapia (*Oreochromis niloticus*) diet. *J Anim Plant Sci*. 2020; 30(2):364–370.
32. Azaza MS, Kammoun W, Abdelmouleh A, Kraïem MM. Growth performance, feed utilization, and body composition of Nile tilapia (*Oreochromis niloticus* L.) fed with differently heated soybean-meal-based diets. *Aquac Int*. 2009; 17(6):507–521.
33. El-Saidy DMSD. Effect of using okara meal, a by-product from soymilk production as a dietary protein source for Nile tilapia (*Oreochromis niloticus* L.) mono-sex males. *Aquac Nutr*. 2011;17(4):380–386.
34. Lin S, Luo L. Effects of different levels of soybean meal inclusion in replacement for fish meal on growth, digestive enzymes and transaminase activities in practical diets for juvenile tilapia, *Oreochromis niloticus* × *O. aureus*. *Anim Feed Sci Technol*. 2011; 168(1):80–87.
35. Koumi AR, Koffi KM, Atsé BC, Kouame LP. Growth, feed efficiency and carcass mineral composition of *Heterobranchius longifilis*, *Oreochromis niloticus* and *Sarotherodon melanotheron* juveniles fed different dietary levels of soybean meal-based diets. *African J Biotechnol*. 2011; 10(66):14990–14998.
36. Koumi AR, Atse BC, Kouame LP. Utilization of soya protein as an alternative protein source in *Oreochromis niloticus* diet: Growth performance, feed utilization, proximate composition and organoleptic characteristics. *African J Biotechnol*. 2009; 8(1):91–97.
37. Obirikorang KA, Gyamfi S, Goode ME, Amisah S, Edziyie RE, Quagrainie K, et al. Effect of soybean meal diets on the growth performance, ammonia excretion rates, gut histology and feed cost of Nile tilapia (*Oreochromis niloticus*) fry. *Aquac Res*. 2020 Sep 1; 51(9):3520–3532.
38. Sharda S, Saini V. Replacement of fishmeal with soybean meal in Nile tilapia (*Oreochromis niloticus*) Diet. *J Entomol Zool Stud*. 2017; 5(4):845–849.
39. Yigit N. O. and Demir T. Use of Fermented Soybean Meal with Whey as a Protein Source for Feeding Juvenile Tilapia (*Oreochromis niloticus*). *Isr J Aquac - Bamidgeh*. 2010;(January):1–7
40. Ashour M, Mabrouk MM, Ayoub HF, El-Feky MMMM, Zaki SZ, Hoseinifar SH, et al. Effect of dietary seaweed extract supplementation on growth, feed utilization, hematological indices, and non-specific immunity of Nile Tilapia, *Oreochromis niloticus* challenged with *Aeromonas hydrophila*. *J Appl Phycol*. 2020; 32(5):3467–3479.
41. Eissa HA, Hegazi MM, Elmor ME, Sharawy ZZ. Effects of partial substitution of fish meal with different levels of marine macroalgae species on growth indices and RNA/DNA ratio of hybrid red tilapia. *Egypt J Aquat Biol Fish*. 2021; 25(3):395–409.
42. Saleh NE, Shalaby SM, Sakr EM, Elmonem AIA, Michael FR. Effect of dietary inclusion of *Ulva fasciata* on red hybrid tilapia growth and carcass composition. *J Appl Aquac*. 2014; 26(3):197–207.
43. Saleh HHE sayed. Can Artificial Feed be Replace by Fresh Macro Algae (*Enteromorpha flaxusa*) in Hybrid Red Tilapia (*Oreochromis mossambicus* × *Oreochromis niloticus*) Juvenile Nutrition? *J Oceanogr Mar Res*. 2020;8(2):1–7.
44. EL-Sayed Hussein EM, Ebtehal EL-Sayed Hussein CM. Effect of seaweed supplemented diets on Nile tilapia, *Oreochromis niloticus* performance. *Int J Fish Aquat Stud*. 2017; 5(2):205–210.
45. Marinho G, Nunes C, Sousa-Pinto I, Pereira R, Rema P, Valente LMP. The IMTA-cultivated Chlorophyta *Ulva* spp. as a sustainable ingredient in Nile tilapia (*Oreochromis niloticus*) diets. *J Appl Phycol*. 2013; 25(5):1359–1367.
46. Radwan M, El-Sharkawy MA, Negm MA, Mohammadein A, Malki JSA, Al-Thomali AW, et al. Dual effect of dietary seaweed of extract nanoparticles (GNS) with bionanocomposite cellulose acetate membranes (CA/bio-AgNps) on growth performance and health status of the Nile tilapia (*Oreochromis niloticus*): Specification on feed utilization, immune s. *Front Mar Sci*. 2022; 9(October):1–18.
47. Salem MES, Abdel-Ghany HM, Almishferi HM. Role of dietary *Laurencia obtusa* in enhancing growth, blood indices, and hypoxia resistance of red tilapia (*Oreochromis niloticus* × *O. mossambicus*). *J Appl Phycol*. 2021; 33(4):2617–2628.
48. Silva DM, Valente LMP, Sousa-Pinto I, Pereira R, Pires MA, Seixas F, et al. Evaluation of IMTA-produced seaweeds (Gracilaria, Porphyra, and Ulva) as dietary ingredients in Nile tilapia, *Oreochromis*

- niloticus* L., juveniles. Effects on growth performance and gut histology. J Appl Phycol. 2015; 27(4):1671–1680.
49. Stadtlander T, Khalil WKB, Focken U, Becker K. Effects of low and medium levels of red alga Nori (*Porphyra yezoensis* Ueda) in the diets on growth, feed utilization and metabolism in intensively fed Nile tilapia, *Oreochromis niloticus* (L.). Aquac Nutr. 2013; 19(1):64–73.
 50. Khalafalla MM and El-Hais A-e.M.A. Evaluation of Seaweeds *Ulva rigida* and *Pterocladia capillacea* as Dietary Supplements in Nile Tilapia Fingerlings. J Aquac Res Dev. 2015;06(03):1–5.
 51. Bakke-McKellep AM, Penn MH, Salas PM, Refstie S, Sperstad S, Landsverk T, et al. Effects of dietary soyabean meal, inulin and oxytetracycline on intestinal microbiota and epithelial cell stress, apoptosis and proliferation in the teleost Atlantic salmon (*Salmo salar* L.). Br J Nutr. 2007; 97(4):699–713.
 52. Cerezo IM, Fumanal M, Tapia-Paniagua ST, Bautista R, Anguís V, Fernández-Díaz C, et al. Solea senegalensis Bacterial Intestinal Microbiota Is Affected by Low Dietary Inclusion of *Ulva ohnoi*. Front Microbiol. 2022; 12(February):1–13.
 53. Ferreira M, Abdelhafiz YAA, Abreu H, Silva J, Valente LMP, Viswanath K. *Gracilaria gracilis* and *Nannochloropsis oceanica*, singly or in combination, in diets alter the intestinal microbiota of European seabass (*Dicentrarchus labrax*). Front. Mar. Sci. 2022; 9:1–16.
 54. Thépot V, Campbell AH, Rimmer MA, Jelocnik M, Johnston C, Evans B, et al. Dietary inclusion of the red seaweed *Asparagopsis taxiformis* boosts production, stimulates immune response and modulates gut microbiota in Atlantic salmon, *Salmo salar*. Aquaculture. 2022; 546:1–13.
 55. Ringø E, Sperstad S, Kraugerud OF, Kroghdahl A. Use of 16S rRNA gene sequencing analysis to characterize culturable intestinal bacteria in Atlantic salmon (*Salmo salar*) fed diets with cellulose or non-starch polysaccharides from soy. Aquac Res. 2008; 39(10):1087–1100.
 56. Rico RM, Tejedor-Junco MT, Tapia-Paniagua ST, Alarcón FJ, Mancera JM, López-Figueroa F, et al. Influence of the dietary inclusion of *Gracilaria cornea* and *Ulva rigida* on the biodiversity of the intestinal microbiota of *Sparus aurata* juveniles. Aquac Int. 2016; 24:965–984.
 57. Reveco FE, Øverland M, Romarheim OH, Mydland LT. Intestinal bacterial community structure differs between healthy and inflamed intestines in Atlantic salmon (*Salmo salar* L.). Aquaculture. 2014;420–421:262–269.
 58. Navarrete P, Fuentes P, De la Fuente L, Barros L, Magne F, Opazo R, et al. Short-term effects of dietary soybean meal and lactic acid bacteria on the intestinal morphology and microbiota of Atlantic salmon (*Salmo salar*). Aquac Nutr. 2013; 19(5):827–836.
 59. Merrifield DL, Dimitroglou A, Bradley G, Baker RTM, Davies SJ. Soybean meal alters autochthonous microbial populations, microvilli morphology and compromises intestinal enterocyte integrity of rainbow trout, *Oncorhynchus mykiss* (Walbaum). J Fish Dis. 2009 Sep 1; 32(9):755–766.
 60. Liu X, Han B, Xu J, Zhu J, Hu J, Wan W, et al. Replacement of fishmeal with soybean meal affects the growth performance, digestive enzymes, intestinal microbiota and immunity of *Carassius auratus gibelio* × *Cyprinus carpio*. Aquac Reports. 2020; 18:1–9.
 61. Keating C, Bolton-Warberg M, Hinchcliffe J, Davies R, Whelan S, Wan AHL, et al. Temporal changes in the gut microbiota in farmed Atlantic cod (*Gadus morhua*) outweigh the response to diet supplementation with macroalgae. Anim Microbiome. 2021; 3:1–21.
 62. Abdala-Díaz RT, García-Márquez J, Rico RM, Gómez-Pinchetti JL, Mancera JM, Figueroa FL, et al. Effects of a short pulse administration of *Ulva rigida* on innate immune response and intestinal microbiota in *Sparus aurata* juveniles. Aquac Res. 202; 52(7):3038–3051.
 63. Heikkinen J, Vielma J, Kemiläinen O, Tirola M, Eskelinen P, Kiuru T, et al. Effects of soybean meal based diet on growth performance, gut histopathology and intestinal microbiota of juvenile rainbow trout (*Oncorhynchus mykiss*). Aquaculture. 2006; 261(1):259–268.
 64. Gonçalves AT, Simões M, Costa C, Passos R, Baptista T. Modulatory effect of *Gracilaria gracilis* on European seabass gut microbiota community and its functionality. Sci Rep. 2022; 12(1):1–16.
 65. Wang Z, Yang L, Zhang C, Li X, Wang L, Lu K, et al. Substituting dietary fishmeal with soybean meal isolate influences hepatic lipid metabolism and gut microbiota in spotted seabass (*Lateolabrax maculatus*). Isr J Aquac - Bamidgah. 2021; 73:1–13.
 66. Tapia-Paniagua ST, Fumanal M, Anguís V, Fernández-Díaz C, Alarcón FJ, Moriñigo MA, et al. Modulation of intestinal microbiota in *Solea senegalensis* fed low dietary level of *Ulva ohnoi*. Front Microbiol. 2019; 10:1–16.
 67. Liu T, Han T, Wang J, Liu T, Bian P, Wang Y, et al. Effects of replacing fish meal with soybean meal on growth performance, feed utilization and physiological status of juvenile redlip mullet *Liza haematocheila*. Aquac Reports. 2021; 20:1–10.

68. Howlader S, Sumi KR, Sarkar S, Billah SM, Ali ML, Howlader J, et al. Effects of dietary replacement of fish meal by soybean meal on growth, feed utilization, and health condition of stinging catfish, *Heteropneustes fossilis*. Saudi J Biol Sci. 2023; 30(3):1–7.
69. Ye H, Xu M, Liu Q, Sun Z, Zou C, Chen L, et al. Effects of replacing fish meal with soybean meal on growth performance, feed utilization and physiological status of juvenile obscure puffer, *Takifugu obscurus*. Comp Biochem Physiol Part - C Toxicol Pharmacol. 2019; 216(November):75–81.
70. Abdel-Warith AA, Younis EM, Al-Asghar NA, Mahboob S. Effect of replacing fish meal by full fat soybean meal on growth performance, feed utilization and gastrointestinal enzymes in diets for african catfish *Clarias gariepinus*. Brazilian J Biol. 2020; 80(3):535–543.
71. Bonaldo A, Roem AJ, Pecchini A, Grilli E, Gatta PP. Influence of dietary soybean meal levels on growth, feed utilization and gut histology of Egyptian sole (*Solea aegyptiaca*) juveniles. Aquaculture. 2006; 261(2):580–586.
72. Kokou F, Rigos G, Henry M, Kentouri M, Alexis M. Growth performance, feed utilization and non-specific immune response of gilthead sea bream (*Sparus aurata* L.) fed graded levels of a bioprocessed soybean meal. Aquaculture. 2012;364–365:74–81.
73. Natify W, Droussi M, Berday N, Araba A, Benabid M. Effect of the seaweed *Ulva lactuca* as a feed additive on growth performance, feed utilization and body composition of Nile tilapia (*Oreochromis niloticus* L.). Int J Agron Agric Res. 2015; 7(3):85–92.
74. Sulaeman HA, Zainuddin, Laining A. Evaluation of seaweed, *Ulva lactuca* as a fresh diet for nursery stage of golden rabbitfish, *Siganus guttatus*. IOP Conf Ser Earth Environ Sci. 2022; 1119(1):1–11.
75. Madibana MJ, Mlambo V, Lewis B, Fouché C. Effect of graded levels of dietary seaweed (*Ulva* sp.) on growth, hematological and serum biochemical parameters in dusky kob, *Argyrosomus japonicus*, sciaenidae. Egypt J Aquat Res. 2017; 43(3):249–254.
76. Seyedalhosseini SH, Salati AP, Mozanzadeh MT, Parrish CC, Shahriari A. Effects of dietary seaweeds (*Gracilaria* spp. and *Sargassum* spp.) on growth, feed utilization, and resistance to acute hypoxia stress in juvenile Asian seabass (*Lates calcarifer*). Aquac Reports. 2023; 31(June):1–6.
77. Sinha AK, Kumar V, Makkar HPS, De Boeck G, Becker K. Non-starch polysaccharides and their role in fish nutrition - A review. Food Chem. 2011; 127(4):1409–1426.
78. Francis G, Makkar HPS, Becker K. Antinutritional factors present in plant-derived alternate fish feed ingredients and their effects in fish. Vol. 199, Aquaculture. 2001. 197–227 p.
79. Kumar V, Sakawat Hossain M, A. Ragaza J, Rubio Benito M. The Potential Impacts of Soy Protein on Fish Gut Health. Soybean Hum Consum Anim Feed. 2020;(September):91–115.
80. Raja K, Kadirvel V, Subramaniyan T. Seaweeds, an aquatic plant-based protein for sustainable nutrition - A review. Futur Foods. 2022; 5(March):1–12.
81. Yang B, Jiang WD, Wu P, Liu Y, Zeng YY, Jiang J, et al. Soybean isoflavones improve the health benefits, flavour quality indicators and physical properties of grass carp (*Ctenopharygodon idella*). PLoS One. 2019; 14(1):1–21.
82. Hoseinifar SH, Jahazi MA, Nikdehghan N, Van Doan H, Volpe MG, Paolucci M. Effects of dietary polyphenols from agricultural by-products on mucosal and humoral immune and antioxidant responses of convict cichlid (*Amatitlania nigrofasciata*). Aquaculture. 2020; 517:1–31.
83. Jahazi MA, Hoseinifar SH, Jafari V, Hajimoradloo A, Van Doan H, Paolucci M. Dietary supplementation of polyphenols positively affects the innate immune response, oxidative status, and growth performance of common carp, *Cyprinus carpio* L. Aquaculture. 2020; 517:1–33.
84. Siddik MAB, Francis P, Rohani MF, Azam MS, Mock TS, Francis DS. Seaweed and Seaweed-Based Functional Metabolites as Potential Modulators of Growth, Immune and Antioxidant Responses, and Gut Microbiota in Fish. Antioxidants. 2023; 12(12):1–34. <https://doi.org/10.3390/antiox12122066> PMID: 38136186
85. Wassef EA, El-Sayed AFM, Sakr EM. Pterocladia (Rhodophyta) and Ulva (Chlorophyta) as feed supplements for European seabass, *Dicentrarchus labrax* L., fry. J Appl Phycol. 2013; 25(5):1369–1376.
86. Martínez-Antequera FP, Martos-Sitcha JA, Reyna JM, Moyano FJ. Evaluation of the inclusion of the green seaweed *Ulva ohnoi* as an ingredient in feeds for gilthead sea bream (*Sparus aurata*) and european sea bass (*Dicentrarchus labrax*). Animals. 2021; 11(6):1–19.
87. Azaza MS, Mensi F, Ksouri J, Dhraief MN, Brini B, Abdelmouleh A, et al. Growth of Nile tilapia (*Oreochromis niloticus* L.) fed with diets containing graded levels of green algae *Ulva rigida* reared in geothermal waters of southern Tunisia. J Appl Ichthyol. 2008; 24(2):202–207.
88. Rahmdel KJ, Noveirian HA, Falahatkar B, Lashkan AB. Effects of replacing fish meal with sunflower meal on growth performance, body composition, hematological and biochemical indices of common carp (*Cyprinus carpio*) fingerlings. Fish Aquat Life. 2018; 26(2):121–129.

89. Güroy D, Güroy B, Merrifield DL, Tekinay AA, Davies SJ, Şahin I. Effects of fish oil and partial fish meal substitution with oilseed oils and meals on growth performance, nutrient utilization and health of the rainbow trout *Oncorhynchus mykiss*. *Aquac Int*. 2012; 20(3):481–497.
90. Elabd H, Soror E, El-Asely A, El-Gawad EA, Abbass A. Dietary supplementation of Moringa leaf meal for Nile tilapia *Oreochromis niloticus*: Effect on growth and stress indices. *Egypt J Aquat Res*. 2019; 45(3):265–271.
91. Yue YR, Zhou QC. Effect of replacing soybean meal with cottonseed meal on growth, feed utilization, and hematological indexes for juvenile hybrid tilapia, *Oreochromis niloticus* × *O. aureus*. *Aquaculture*. 2008; 284(1):185–189.
92. Ibrahim A, Abouelella S, Adam H, Hassan N, Abdallah R. Effect of Azolla pinnata on growth performance and survival rate of fingerlings of grass carp fish *Ctenopharyngodon idellus* (Valenciennes, 1844). *Aswan Univ J Environ Stud*. 2021; 2(4):249–258.
93. Moyo NAG, Rapatsa-Malatji MM. A review and meta-analysis of selected plant protein sources as a replacement of fishmeal in the diet of tilapias. *Ann Anim Sci*. 2023; 23(3):681–690.
94. Xia Z, Zhu M, Zhang Y. Effects of the probiotic *Arthrobacter* sp. CW9 on the survival and immune status of white shrimp (*Penaeus vannamei*). *Lett Appl Microbiol*. 2014; 58(1):60–64.
95. Wuertz S, Beça F, Kreuz E, Wanka KM, Azeredo R, Machado M, et al. Two Probiotic Candidates of the Genus *Psychrobacter* Modulate the Immune Response and Disease Resistance after Experimental Infection in Turbot (*Scophthalmus maximus*, Linnaeus 1758). *Fishes*. 2023; 8(3):12–14.
96. Moroni F, Naya-Català F, Piazzon MC, Rimoldi S, Caldach-Giner J, Giardini A, et al. The Effects of Nisin-Producing *Lactococcus lactis* Strain Used as Probiotic on Gilthead Sea Bream (*Sparus aurata*) Growth, Gut Microbiota, and Transcriptional Response. *Front Mar Sci*. 2021; 8(April):1–18.
97. Vasyliuk OM, Skrotskyi SO, Khomenko LA, Babich T V. Probiotics Based on Lactic Acid Bacteria for Aquaculture. *Mikrobiol Zh*. 2023; 85(2):75–92.
98. Hu Y, Zhang J, Xue J, Chu W, Hu Y. Effects of dietary soy isoflavone and soy saponin on growth performance, intestinal structure, intestinal immunity and gut microbiota community on rice field eel (*Monopterus albus*). *Aquaculture*. 2021; 537:1–11.
99. Zhou C, Lin H, Ge X, Niu J, Wang J, Wang Y, et al. The Effects of dietary soybean isoflavones on growth, innate immune responses, hepatic antioxidant abilities and disease resistance of juvenile golden pompano *Trachinotus ovatus*. *Fish Shellfish Immunol* [Internet]. 2015; 43(1):158–166.
100. Chojnacka K. Biologically Active Compounds in Seaweed Extracts - the Prospects for the Application. *Open Conf Proc J*. 2012; 3(1):20–28.
101. Hentati F, Tounsi L, Djomdi D, Pierre G, Delattre C, Ursu AV, et al. Bioactive polysaccharides from seaweeds. *Molecules*. 2020; 25(14):1–29. <https://doi.org/10.3390/molecules25143152> PMID: 32660153
102. Kumar LRG, Paul PT, Anas KK, Tejpal CS, Chatterjee NS, Anupama TK, et al. Phlorotannins—bioactivity and extraction perspectives. *J Appl Phycol*. 2022; 34(4):2173–2185. <https://doi.org/10.1007/s10811-022-02749-4> PMID: 35601997
103. Ramnani P, Chittarari R, Tuohy K, Grant J, Hotchkiss S, Philp K, et al. In vitro fermentation and prebiotic potential of novel low molecular weight polysaccharides derived from agar and alginate seaweeds. *Anaerobe*. 2012; 18(1):1–6.
104. Alvanou M V, Feidantsis K, Staikou A, Apostolidis AP, Michaelidis B, Giantsis IA. Probiotics, Prebiotics, and Synbiotics Utilization in Crayfish Aquaculture and Factors Affecting Gut Microbiota. *Microorganisms*. 2023; 11(5):1–30. <https://doi.org/10.3390/microorganisms11051232> PMID: 37317206
105. Eom SH, Kim YM, Kim SK. Antimicrobial effect of phlorotannins from marine brown algae. *Food Chem Toxicol*. 2012; 50(9):3251–3255. <https://doi.org/10.1016/j.fct.2012.06.028> PMID: 22735502
106. Ford L, Stratakis AC, Theodoridou K, Dick JTA, Sheldrake GN, Linton M, et al. Polyphenols from Brown Seaweeds as a Potential Antimicrobial Agent in Animal Feeds. *ACS Omega*. 2020; 5(16):9093–9103. <https://doi.org/10.1021/acsomega.9b03687> PMID: 32363261
107. Manso T, Lores M, de Miguel T. Antimicrobial Activity of Polyphenols and Natural Polyphenolic Extracts on Clinical Isolates. *Antibiotics*. 2022; 11(1):1–18.
108. Pradhan B, Nayak R, Bhuyan PP, Patra S, Behera C, Sahoo S, et al. Algal Phlorotannins as Novel Antibacterial Agents with Reference to the Antioxidant Modulation: Current Advances and Future Directions. *Mar Drugs*. 2022; 20(6):1–15.
109. Zhang XY, Zeng XN, Liu S, Wu F, Li YY, Pan Q. Effects of dietary four different woody forages on gut microbiota of Nile tilapia (*Oreochromis niloticus*). *Aquac Res*. 2021; 52(4):1733–1742.
110. Wu F, Chen B, Liu S, Xia X, Gao L, Zhang X, et al. Effects of woody forages on biodiversity and bioactivity of aerobic culturable gut bacteria of tilapia (*Oreochromis niloticus*). *PLoS One*. 2020; 15(7):1–17.