

**PREVALENCE OF AFLATOXIN M₁ IN PASTEURIZED AND ULTRA-HIGH
TEMPERATURE (UHT) MILK MARKETED IN DAR ES SALAAM, TANZANIA**

HILDA FREDY MWAKOSYA

**A DISSERTATION SUBMITTED IN PARTIAL FULFILMENT
OF THE REQUIREMENTS FOR THE DEGREE OF MASTER OF SCIENCE
IN FOOD QUALITY AND SAFETY ASSURANCE OF SOKOINE UNIVERSITY
OF AGRICULTURE, MOROGORO, TANZANIA**

EXTENDED ABSTRACT

A survey was conducted in year 2020/2021 to establish the levels of aflatoxin M₁ (AFM₁) in pasteurized and ultra-heat-treated milk (n=118) and awareness on mycotoxins by milk processors in Kinondoni, Temeke, Ubungo, Ilala and Kigamboni district of Dar es Salaam, the commercial capital of Tanzania. The levels of AFM₁ in pasteurized milk (n=75) and UHT milk (n=43) samples were determined by using immuno-affinity high performance liquid chromatography. AFM₁ was detected in 97% (115/118) of the heat-treated samples. Pasteurized milk and UHT milk samples were contaminated by 96% (72/75) and 100% (43/43), respectively. About 82% of the contaminated pasteurized and UHT had aflatoxin M₁ above the EU acceptable levels (0.05 µg/L) however none of the contaminated pasteurized and UHT milk sample exceed Codex limits of 0.5 µg/L. The observed contamination levels of AFM₁ in heated milk could pose a serious public health problem. Therefore, best practices including regular monitoring of AFM₁ levels in milk and milk products are crucial to protect consumers. Awareness of aflatoxin contamination of milk was assessed by using a cross-sectional descriptive statistic involving 30 milk processors. Data was collected using a structured questionnaire and statistically analysed using Statistical Package for Social Sciences (IBM SPSS® Version 27 (2020)). Descriptive statistics was used to determine frequencies and percentages of social demographic, knowledge, handling and feeding practices of lactating cow. Cross tabulation was used to determine relationship between knowledge on aflatoxins with age and education level of the respondents. The majority of the respondents (83.3%) were aware of aflatoxins and none (0.0%) of the respondents were aware that milk and milk products could be contaminated with aflatoxins. It was also observed that the cattle feeding practices were poor and were a major reason for AFM₁ contamination of milk. None of the respondents were aware that feeding lactating cow with mouldy feeds could results into AFM₁ contamination of milk. It was observed that, AFM₁ analysis was not carried out in raw

milk before processing in order to control AFM₁ contamination of milk and milk products. This could be due to lack of knowledge and techniques for detection and analysis of aflatoxins. This study recommended that creation of awareness of aflatoxins and use of best practices along the milk value chain was crucial in order to enhance the safety of consumers.

DECLARATION

I, Hilda Mwakosya do hereby declare to the Senate of Sokoine University of Agriculture that this dissertation is my own original work done within the period of registration and that it has neither been submitted nor being concurrently submitted in any other institution for a degree award.

Hilda Mwakosya

(MSc. Food Quality and Safety Assurance)

Date

The above declaration is confirmed;

Prof. Jovin K. Mugula

(Supervisor)

Date

COPYRIGHT

No part of this dissertation may be reproduced, stored in any retrieval system, or transmitted in any form or by any means without prior written permission of the author or the Sokoine University of Agriculture in that behalf.

ACKNOWLEDGEMENTS

I wish to thank the Tanzania Bureau of Standards (TBS) for giving me the opportunity, study leave and financially support to pursue this master's degree programme.

I wish to express my gratitude to my supervisor Prof. Jovin K. Mugula for his guidance, continuous encouragement and support from the proposal write up to submission of the dissertation.

I am thankful to the food laboratory staff in the Directorate of Testing and Calibration at Tanzania Bureau of Standards (TBS), particularly Mr. Roman M. Fortunatus and Mr. Wiseborn Makaka for technical support during my laboratory work. I also acknowledge the assistance from the Tanzania Dairy Board (TDB) for locating milk processors in Dar es salaam. I am grateful to milk processors in Dar es salaam for their willingness to participate in this study.

I am thankful for the support given by my colleagues in Msc. Food Quality and Safety Assurance students of the year 2019 to 2021 particularly Catherine Akwilapo, Joseph Samwel, Saimon Pela, Asha Ndwata and Rose Jumbe in the Department of Food Technology, Nutrition and Consumer Sciences at SUA.

Last but not least, I wish to express my appreciation to my family and my friends for inspiration, encouragement, love and support. I am especially obligated to my husband Mr. Daniel Amulike Aron and my children Hellen, Allen and Grayson for their patience and understanding during my absence from home.

DEDICATION

This dissertation is lovingly dedicated to my husband, Mr. Daniel Amulike Aron for his support, encouragement and constant love and to my children Hellen, Allen and Grayson for their love and patience during my absence.

TABLE OF CONTENTS

EXTENDED ABSTRACT.....	ii
DECLARATION.....	iv
COPYRIGHT.....	v
ACKNOWLEDGEMENTS.....	vi
DEDICATION.....	vii
TABLE OF CONTENTS.....	viii
LIST OF TABLES.....	xi
LIST OF APPENDICES.....	xii
LIST OF ABBREVIATION AND ACRONYMS.....	xiii
CHAPTER ONE.....	1
1.0 INTRODUCTION.....	1
1.1 Background Information.....	1
1.1.1 Milk production in Tanzania.....	1
1.1.2 Aflatoxin M ₁ in heat treated milk.....	1
1.1.3 Ill-health effects of aflatoxin M ₁	3
1.1.4 Prevalence of raw and heat-treated milk with aflatoxins.....	3
1.2 Justification.....	4
1.3 Objectives.....	5
1.3.1 Main objective.....	5
1.3.2 Specific objectives.....	5
References.....	6

CHAPTER TWO.....	10
2.0 PREVALENCE OF AFLATOXIN M₁ IN PASTEURIZED AND ULTRA-HIGH TEMPERATURE (UHT) MILK MARKETED IN DAR ES SALAAM, TANZANIA.....	10
CHAPTER THREE.....	17
3.0 KNOWLEDGE, HANDLING AND FEEDING PRACTICES ASSOCIATED WITH AFLATOXIN M₁ CONTAMINATION OF MILK IN DAR ES SALAAM, TANZANIA.....	17
3.1 Abstract.....	17
3.2 Introduction.....	18
3.3 Materials and Methods.....	20
3.3.1 Study design.....	20
3.3.2 Study population.....	21
3.3.3 Sampling techniques and procedures.....	21
3.3.4 Data collection tools.....	21
3.3.5 Statistical analysis.....	22
3.4 Results and discussion.....	22
3.4.1 Demographic characteristics of the respondents.....	22
3.4.2 Knowledge on aflatoxins contamination.....	23
3.4.3 Milk handling practices.....	26
3.4.4 Feeding practices of lactating cow by the study respondents.....	27
3.4.5 Relationship between knowledge of aflatoxin with age and level of education.....	30
3.5 Conclusion.....	32
3.6 Acknowledgements.....	32

References.....33

CHAPTER FOUR.....39

4.0 OVERALL CONCLUSION AND RECOMMENDATIONS.....39

4.1 Conclusions.....39

4.2 Recommendations.....40

APPENDICES.....42

Table 3.1:	Demographic characteristics of the study respondents.....	23
Table 3.2:	Knowledge on aflatoxins by the study respondents.....	26
Table 3. 3:	Handling practices of milk by the study respondents.....	27
Table 3.4:	Feeding practices of lactating cow by the study respondents.....	29
Table 3.5:	Relationship between knowledge on aflatoxins by education level of the study respondents.....	30
Table 3.6:	Relationship between knowledge of aflatoxins with age of the study respondents.....	31

Appendix 1: Photos showing poor storage practices of cattle feeds for feeding cow.....42

Appendix 2: Questionnaire to assess knowledge and practices on aflatoxin
contamination in dairy products to milk processors.....43

LIST OF ABBREVIATION AND ACRONYMS

AFB₁ Aflatoxin B₁

AFB ₂	Aflatoxin B ₂
AFG ₁	Aflatoxin G ₁
AFG ₂	Aflatoxin G ₂
AFM ₁	Aflatoxin M ₁
AFM ₂	Aflatoxin M ₂
CAC	Codex Alimentarius Commission
EU	European Union
FAO	Food and Agriculture Organization
FLD	Fluorescence Detector
GAP	Good Agricultural Practices
GDP	Gross Domestic Product
HPLC	High Performance Liquid Chromatography
IARC	International Agency for Research on Cancer
LOD	Limit of Detection
LOQ	Limit of Quantification
MCT	Medium-chain triglycerides
µg/L	Microgram per Liter (ppb)
RPM	Revolution per Minutes
SPSS	Statistical Package for Social Sciences
SUA	Sokoine University of Agriculture
TBS	Tanzania Bureau of Standards
UHT	Ultra High Temperature
WHO	World Health Organization

CHAPTER ONE

1.0 INTRODUCTION

1.1 Background Information

1.1.1 Milk production in Tanzania

Tanzania has the third largest livestock population in Africa comprising of 25 million cattle out of which 98% are indigenous breeds (FAO, 2020). In year 2018 about 2.09 billion litres of raw milk were produced in Tanzania and milk production contributes to income, food security, nutrition and household livelihood (FAOSTAT, 2020). The sector contributes to 7.4% of total national GDP, however the annual growth rate (2.2 %) of the sector is low (FAO, 2020).

The dairy production sector in Tanzania is divided into two subsectors: traditional and improved dairy subsectors (Nell *et al.*, 2014). Traditional subsector is the most dominant and dominated by Tanzanian Shorthorn Zebu (TSHZ) cattle with production of 70% milk of which 90% is consumed at home and 10% contributes to commercial sector (Munyaneza *et al.*, 2019; URT, 2017). The improved dairy system subsector is dominated by smallholder's dairy farmers who keep mainly improved dairy cattle and contributes about 30% of all milk produced and marketed in Tanzania (URT, 2017).

1.1.2 Aflatoxin M₁ in heat treated milk

Raw milk is a valuable nutritious food. However, it is highly perishable (has a short shelf-life) and an excellent medium for the growth of microorganisms, particularly bacterial pathogens that can cause spoilage and diseases to consumers (FAO, 2021). Heat treatment of milk such as pasteurization and UHT (ultra-high temperature) allows the preservation of milk and helps to reduce food-borne illness (Melini *et al.*, 2017).

According to FAO/WHO (1982), pasteurization is defined as a heat treatment process applied to a product such as milk with the objective of minimizing numbers of harmful micro-organisms to a level at which they do not constitute a significant health hazard with minimal chemical, physical and organoleptic changes in the product. It also extends the storage time for some products by reducing the number of spoilage micro-organisms in the product (FAO/WHO,1982). Codex Alimentarius (2004), defined UHT (ultra-high temperature) treatment of milk and liquid milk products as the application of heat to a continuously flowing product using such high temperatures for short time that renders the product commercially sterile at the time of processing. When UHT treatment is combined with aseptic packaging, it results in a commercially sterile product at the heating range of 135 to 150 °C for 1 s up to 4 s (Melini *et al.*,2017).

Aflatoxins are amongst the most poisonous mycotoxins and are produced by *Aspergillus flavus*, *Aspergillus parasiticus* and *Aspergillus nomius* fungi found in soil and that can grow in plant, human food products and feeds (WHO, 2018). The most important aflatoxins in order of toxicity are B₁, B₂, G₁, and G₂ (Ismail *et al.*, 2018; Tahira *et al.*, 2019). Aflatoxins may also be found in the milk of animals that are fed contaminated feed, in the form of aflatoxin M₁ (Langat *et al.*, 2016).

Aflatoxin M₁ (AFM₁) is the principal metabolite of aflatoxin B₁ (AFB₁), and it is formed in the liver (Langat *et al.*, 2016). Once lactating cow consume contaminated feeds with aflatoxin B₁, it is absorbed into the gastro intestinal tract and biotransformation occurs in the liver by cytochrome P₄₅₀ enzymes to form a 4-hydroxy metabolite known as aflatoxin M₁ a compound soluble in water and therefore it is easily excreted in milk during milking and appears within 12 hours of administration of contaminated feeds (Daou *et al.*,2020; Tahira *et al.*, 2019).

AFM₁ is a heat stable compound that can survive heat treatment such as pasteurization, UHT technique and autoclaving but also AFM₁ may be reduced but not completely destroyed by heat treatments (Mahmoodi *et al.*, 2019; Tahira *et al.*, 2019).

1.1.3 Ill-health effects of aflatoxin M₁

Aflatoxin M₁ is a hepato-carcinogen, classified as a group 1 carcinogen by International Agency for Research on Cancer (IARC Monograph, 2018). Exposure of humans to aflatoxin M₁ leads to several health-related problems including acute and chronic aflatoxicosis; it exerts its negative effect on health through binding to nucleic acid causing DNA damage and eventually leading to hepatotoxicity, carcinogenicity, immune suppression, cirrhosis and stunted growth in children (Singh *et al.*, 2021).

1.1.4 Prevalence of raw and heat-treated milk with aflatoxins

Prevalence of milk with aflatoxin M₁ is due to consumption of feeds contaminated with aflatoxin B₁, which eventually contaminate milk (Langat *et al.*, 2016). Many studies have been conducted on aflatoxin M₁ contamination in raw and heat treated milk, with findings revealing levels above the Codex Alimentarius and EU limits from various countries, such as, AFM₁ contamination in pasteurized and UHT milk in Morocco (Mannani *et al.*, 2021); AFM₁ contamination in pasteurized and powdered milk products in Iran (Mahmoodi *et al.*, 2019); AFM₁ contamination in raw, pasteurized, UHT cows' milk and dairy products in Lebanon (Daou, *et al.*, 2020); AFM₁ contamination in milk and milk products in Turkey (Eker *et al.*, 2019); AFM₁ contamination in raw and processed milk in Pakistan (Tahira *et al.*, 2019) and AFM₁ contamination in milk and milk products in Kenya (Langat *et al.*, (2016).

Langat *et al.* (2016) reported that 84.32% (156/185) of the samples of raw milk, processed milk and milk products collected in Bomet county in Kenya were contaminated with aflatoxin M₁ and 43.8% of samples had contamination levels higher than the tolerance limit of 0.05 µg/l recommended by FAO and WHO. In addition, the study indicated that the level of contamination in raw milk was higher (52.0%) than in processed milk (8.6%). Mohammed (2016) reported that 83.8% (31/37) of raw cow milk samples collected randomly from different locations in Singida region, Tanzania was contaminated with AFM₁, with levels exceeding the FAO/WHO of 0.05 ng/ml.

There is scanty information on aflatoxin M₁ contamination in thermal processed cow's milk, particularly, pasteurized and UHT, in Tanzania, as well as knowledge associated with aflatoxin M₁ contamination, as previous studies mostly reported on contamination in raw cow's milk in few regions only. Therefore, the aim of this study was to carry out surveillance on aflatoxin M₁ contamination of pasteurized and UHT milk marketed in Dar es Salaam, the commercial capital of Tanzania.

1.2 Justification

Aflatoxin M₁ is a hepato-carcinogen, classified as a group 1 carcinogen by International Agency for Research on Cancer (IARC Monograph, 2018). Aflatoxin M₁ is a heat-stable and can survive pasteurization, autoclaving and thermal inactivation (Zakaria *et al.*, 2019). The contamination of milk and milk products by aflatoxin M₁ has been reported in various countries such as, AFM₁ contamination in pasteurized and UHT milk in Morocco (Mannani *et al.*, 2021); AFM₁ contamination in pasteurized and powdered milk products in Iran (Mahmoodi *et al.*, 2019); AFM₁ contamination in raw, pasteurized, UHT cows' milk and dairy products in Lebanon (Daou, *et al.*, 2020); AFM₁ contamination in milk and milk products in Turkey (Eker *et al.*, 2019); AFM₁ contamination in raw and processed

milk in Pakistan (Tahira *et al.*, 2019) and AFM₁ contamination in milk and milk products in Kenya (Langat *et al.*, (2016).

Prevalence of aflatoxin M₁ in milk in Tanzania indicated that 92% of raw cow milk retailed in Dar es Salaam city was contaminated with aflatoxin M₁ (Urio *et al.*, 2006) and 83.8% of raw cow milk from households in Singida was contaminated with aflatoxin M₁ (Mohammed *et al.*, 2016). However, there is no reported information on aflatoxin M₁ contamination of pasteurized and ultra-pasteurized (UHT) milk in Tanzania, as well as the awareness of contamination. Thus, the aim of this study was to carry out surveillance on the level of contamination of aflatoxin M₁ in pasteurized and ultra-pasteurized marketed milk in Dar-es-salaam, commercial capital of Tanzania. The results of this study will provide information on level of milk contamination by aflatoxins and contribute to the efforts of food control authorities in developing strategies to ensure public safety. Also, it will serve as a basis for awareness creation for milk value chain stakeholders, including feed millers, milk processors and consumers, on the aflatoxin contamination in pasteurized milk and safety implications.

1.3 Objectives

1.3.1 Main objective

The main objective of this study was to carryout surveillance on aflatoxin M₁ contamination in pasteurized and ultra-pasteurized (UHT) cow milk marketed in Tanzania.

1.3.2 Specific objectives

The specific objectives of this study were to:

- i. determine the prevalence and levels of aflatoxin M₁ in pasteurized and UHT milk marketed in Dar es Salaam.

- ii. assess the knowledge and practices of milk processors on factors associated with aflatoxins contamination in milk and milk products.

The findings of this research work were reported in two manuscripts presented as chapter two and three.

References

- Akhtar, S., Shahzad, M. A., Yoo, S. H., Ismail, A., Hameed, A., Ismail, T., and Riaz, M. (2017). Determination of aflatoxin M₁ and heavy metals in infant formula milk brands available in Pakistani markets. *Korean Journal for Food Science of Animal Resources* 37(1): 79-86.
- Daou, R., Afif, C., Joubrane, K., Khabbaz, L. R., Maroun, R., Ismail, A., and El Khoury, A. (2020). Occurrence of aflatoxin M₁ in raw, pasteurized, UHT cows' milk, and dairy products in Lebanon. *Food Control* 111:1-19.
- Eker, F. Y., Muratoglu, K., and Eser, A. G. (2019). Detection of aflatoxin M₁ in milk and milk products in Turkey. *Environmental Monitoring and Assessment* 191(8): 1-8.
- FAOSTAT (2020). Raw and heat-treated milk production in Tanzania 2018. [<http://www.fao.org/faostat/en/#data/QP>] site visited on 07/06/2021.
- FAO (2021). Gateway to dairy production and products - Milk processing. [<http://www.fao.org/dairy -production-products/processing/en/>] site visited on 21/5/2021.

FAO, (2020): Tanzania at a glance. Food and Agriculture Organization of the United Nations, [<http://www.fao.org/tanzania/fao-in-tanzania/tanzania-at-a-glance/en/>] site visited on 22/9/2020.

International Agency for Research on Cancer. (2018). *Monograph 100F Aflatoxins. 2002*, 225–248. [<https://monographs.iarc.fr/wp-content/uploads/2018/06/mono100F-23.pdf>] site visited on 15/7/2020.

Langat, G., Tetsuhiro, M., Ganoi, T., Matiru, V., and Bii, C. (2016). Aflatoxin M₁ contamination of milk and its products in Bomet County, Kenya. *Advances in Microbiology*, 6(07): 528-536.

Mahmoodi, M., Mazaheri, M., and Talebi Mehrdar, M. (2019). Determination of Aflatoxin M₁ in pasteurized liquid and powdered milk products imported to Iran. *Iranian Journal of Toxicology*, 13(2): 19-23.

Mannani, N., Tabarani, A., El Adlouni, C., and Zinedine, A. (2021). Aflatoxin M₁ in pasteurized and UHT milk marketed in Morocco. *Food Control*, 124: 1-12.

Melini, F., Melini, V., Luziatelli, F., and Ruzzi, M. (2017). Raw and heat-treated milk: From public health risks to nutritional quality. *Beverages* 3(4): 1-33.

Mohammed, S., Munissi, J. J., and Nyandoro, S. S. (2016). Aflatoxin M₁ in raw milk and aflatoxin B₁ in feed from household cows in Singida, Tanzania. *Food Additives and Contaminants: Part B*, 9 (2): 85-90.

- Munyaneza, C., Kurwijila, L. R., Mdoe, N. S., Baltenweck, I., and Twine, E. E. (2019). Identification of appropriate indicators for assessing sustainability of small-holder milk production systems in Tanzania. *Sustainable Production and Consumption* 19: 141-160.
- Nell, A. J., Schiere, H., and Bol, S. (2014). Quick scan dairy sector Tanzania. [<http://www.edepot.wur.nl/334382>] site visited on 21/5/2021.
- Singh, U., Gupta, S., and Gupta, M. (2021). A review study on biological ill effects and health hazards of aflatoxins. *Asian Journal of Advances in Medical Science* 1-8.
- Tahira, I., Sultana, N., Munir, A., Hasan, S. M. and Hanif, N. Q. (2019). Occurrence of Aflatoxin M₁ in raw and processed milk consumed in Pakistan. *Pakistan Journal of Pharmaceutical Sciences*, 32(3): 1097-1101.
- United Republic of Tanzania. (2017). Tanzania livestock master plan (2017/2018 – 2021/2022). Ministry of Livestock and Fisheries.
- Urio, E., Juma, A., Mwanyika, S., Mlingi, N., Ndunguru, G., and Ndossi, G. (2006). The occurrence of aflatoxin M₁ in fresh milk retailed in Dar es Salaam, Tanzania In: *Mycotoxins and phytotoxins*. (Edited by Njapau H, Trujillo S, van Egmond HP and Park DL), Advances in determination, toxicology and exposure management. Wageningen Academic Publishers, Netherlands. pp. 202-207.
- WHO (2018). Aflatoxins – Food Safety Digest. [<http://www.who.int/foodsafety/FSDigestAflatoxinsEN.pdf>]. site visited on 18/5/2021.

WHO (2018). Mycotoxins: World Health Organization. [<https://www.who.int/news-room/fact-sheets/detail/mycotoxins>] site visited on 6/6/2020.

Zakaria, A. M., Amin, Y. A., Khalil, O. S. F., Abdelhiee, E. Y., and Elkamshishi, M. M. (2019). Rapid detection of aflatoxin M₁ residues in market milk in Aswan Province, Egypt and effect of probiotics on its residues concentration. *Journal of Advanced Veterinary and Animal Research*, 6(2): 197–201197.

CHAPTER TWO

PAPER ONE

2.0 PREVALENCE OF AFLATOXIN M₁ IN PASTEURIZED AND ULTRA-HIGH TEMPERATURE (UHT) MILK MARKETED IN DAR ES SALAAM, TANZANIA

H.F. Mwakosya^{1,2} and J. K Mugula²

¹Tanzania Bureau of Standards, P.O. Box 9524, Dar es Salaam, Tanzania.

²Department of Food Technology, Nutrition and Consumer Sciences, Sokoine University of Agriculture, P.O. Box 3006, Morogoro Tanzania.

* Email of the corresponding author: hmwakosya@gmail.com

CHAPTER THREE

3.0 KNOWLEDGE, HANDLING AND FEEDING PRACTICES ASSOCIATED WITH AFLATOXIN M₁ CONTAMINATION OF MILK IN DAR ES SALAAM, TANZANIA.

H.F. Mwakosya^{1,2}, J. K Mugula²

¹Tanzania Bureau of Standards, P.O. Box 9524, Dar es Salaam, Tanzania.

²Department of Food Technology, Nutrition and Consumer Sciences, Sokoine University of Agriculture, P.O. Box 3006, Morogoro Tanzania.

*Email of the corresponding author hmwakosya@gmail.com

3.1 Abstract

This aim of this study was to establish the knowledge, handling and feeding practices associated with aflatoxin M₁ contamination in milk marketed in Dar es Salaam, Tanzania. A cross-sectional descriptive study involving 30 milk processors was conducted. Data was collected using a structured questionnaire and analysed using Statistical Package for Social Sciences (IBM SPSS® Version 27 (2020)). Descriptive statistics was used to determine frequencies and percentages of social demographic, knowledge, handling practices of milk and feeding practices of lactating cow. Cross tabulation was used to determine relationship between knowledge of aflatoxins with age and education level of the respondents. The results indicated that the majority of the respondents (83.3%) were aware of existence of aflatoxins and none (0.0%) of the study respondents were aware of aflatoxin contamination in milk and milk products. It was also observed that feeding practices were poor and could be major contribution for AFM₁ contamination in milk.

Although respondents were aware that aflatoxins caused ill-health effects on humans and animals, but were not aware that feeding lactating cow with mouldy feeds could cause into aflatoxin contamination in milk. It was observed that, there was no AFM₁ analysis carried out in raw milk before processing to control contamination AFM₁ in milk and milk products. Therefore, knowledge creation on aflatoxin contamination and use of best feeding practices along the milk value chain is recommended in order to protect public health (safety).

Keywords: Aflatoxin M₁, Awareness, Handling, Feeding practices, Heat treated Milk, Food safety.

3.2 Introduction

Mycotoxins are toxic secondary metabolites produced mainly by certain species of moulds which contaminate variety of agricultural food stuffs as well as animal derived products such as meat, milk and eggs (Alshannaq *et al.*, 2017; Haque *et al.*, 2020; WHO, 2018). Contamination can occur at any stage in food value chain (Achaglinkame *et al.*, 2017).

It has been reported that about 25% of global food crops are contaminated with mycotoxins to the levels which may pose serious public health effects to human (Eskola *et al.*, 2020). Therefore, contamination of food products with mycotoxin is a global concern because it can result into economic losses, adverse health effects to human and domestic animals and barriers to trade (Magembe *et al.*, 2016). More than 500 mycotoxins have been identified worldwide; however, there are few very toxic mycotoxins that are of public health concern specifically, aflatoxins, ochratoxins, fumonisins, patulin, zearalenone, and trichothecenes such as deoxynivalenol and T-2 toxin (Horky *et al.*, 2018; Palumbo *et al.*, 2020).

Aflatoxins are toxic secondary metabolites produced by toxigenic moulds particularly *Aspergillus* namely *A. flavus*, *A. parasiticus* and *A. nomius*. There are more than 20 molecules of aflatoxins and the most prominent aflatoxins are AFB₁, AFB₂, AFG₁, AFG₂, AFM₁ and AFM₂ (Gurbuz *et al.*, 2019; Ismail *et al.*, 2018). AFM₁ is the 4-hydroxy derivative of AFB₁ and are found in milk and milk products of animals that are fed contaminated feeds with aflatoxin B₁ (Ayofemi, 2020; Lee *et al.*, 2017; Udomkun *et al.*, 2018). The concentration of aflatoxin M₁ in milk is about 0.3 to 6.2% of the concentration of aflatoxin B₁ ingested with the feed and it appears within 12 hours of administration of contaminated feeds, however the concentration level differs depending on factors such as genetics of the animals, seasonal variation, the milking process and the environmental conditions (Ketney *et al.*, 2017; Langat *et al.*, 2016; Vaz *et al.*, 2020). AFM₁ is relatively stable during milk pasteurization, storage, and processing (Mahmoodi *et al.*, 2019). The presence of AFM₁ in milk is a major risk to the public especially children who consume milk as a sole source of diet because of its toxicity and carcinogenicity effect and it has thus been re categorized by International Agency for Research on Cancer (IARC) from Group 2B to Group 1 as a proved carcinogen (IARC, 2012; Li *et al.*, 2017).

Animal feeds such as maize, wheat, paddy (grains, husk or straw), green grass, oil seeds cakes and kitchen wastes are potential source of aflatoxin exposure to animals (Ayo *et al.*, 2018; Choudhary *et al.*, 2020; Tahira *et al.*, 2019). In Tanzania, studies on aflatoxin contamination in cattle feeds was reported in sunflower seeds cake by Mmongoyo *et al.*, (2017) and Mohammed *et al.*, (2016) and in maize by Kamala *et al.* (2016). Therefore, proper handling of these products is required in order to prevent exposure of aflatoxins to animals and in milk and milk products.

Several studies have been conducted on awareness and knowledge on animal feeds and milk contamination with aflatoxins worldwide. In Punjab, dairy farmers were assessed on knowledge and practices to control aflatoxin in the dairy rations (Mir *et al.*, 2020). Moreover, Kagera *et al.*, (2019) conducted a study to assess awareness, knowledge and practices on aflatoxins contamination in milk in urban and peri-urban farmers in Kasarani, Kenya.

In Tanzania, awareness and knowledge on aflatoxins contamination was conducted mainly on maize, groundnuts, animal feeds and spices (Ayo *et al.*, 2018; Magembe *et al.*, 2016; Fundikira *et al.*, 2021). There is limited information on the level of knowledge and practices affecting AFM₁ contamination of milk and milk products by milk processors in Tanzania. Therefore, the aim of this study was to assess the knowledge, handling and feeding practices by milk processors on AFM₁ contamination in milk and milk products in Tanzania. The information obtained from this study will be useful in designing measures for increasing milk processor's awareness towards reduction of AFM₁ exposure to milk and milk products.

3.3 Materials and Methods

3.3.1 Study design

A cross-sectional descriptive study was carried out between December 2020 and January 2021 in Ilala, Kinondoni, Temeke, Ubungo and Kigamboni five districts of Dar es Salaam, the commercial capital of Tanzania. This design enabled collection of data on handling, feeding practices and knowledge on aflatoxins by the study population involved in milk processing.

3.3.2 Study population

The study population was people who were involved in milk processing in five districts of Dar es Salaam. This group was selected because it was involved in milk processing. In addition, these people were more likely to respond to the study questions precisely.

3.3.3 Sampling techniques and procedures

The present study was a cross sectional study design. It examined milk processors at a specific point in time from December 2020 to January 2021. A total of 30 milk processors were interviewed from Kinondoni, Ilala, Ubungo, Temeke and Kigamboni districts in Dar es salaam, eleven milk processors were purposively obtained from list of working dairy industry in Dar es salaam registered by Tanzania Dairy Board (TDB) 2020/2021 and the other 19 milk processors were obtained by snowball technique by face-to-face interview from 11 registered milk processors.

3.3.4 Data collection tools

The data of the study was collected using quantitative methods. An interview administered questionnaire (with closed and open-ended questions) was used to study respondents who consented on participation. This questionnaire was in English and translated to Swahili.

The questionnaire administered to milk processors had forty-three questions that attempted to capture information on knowledge of aflatoxins and issues on milk handling and feeding practices. The questionnaire was pre tested on a random sample of 8 participants (milk processors) at Ilala districts to provide a clear indication on the response and relevance of the questions to the study objectives.

3.3.5 Statistical analysis

The data was analysed using Statistical Package for Social Sciences (IBM SPSS® Version 27 (2020)). The data was coded appropriately and fed into SPSS version 27 to determine the frequencies. The analysis involved descriptive statistics to describe socio-demographic, knowledge, feeding and handling practices, in frequency tables. Cross tabulation (Fisher's exact test at $P < 0.05$) was used to analyse relationship between two variables such as relationship on knowledge of aflatoxins with age of milk processors and relationship on knowledge of aflatoxins with education level of milk processors.

3.4 Results and discussion

3.4.1 Demographic characteristics of the respondents

In this study 30 respondents who were involved in processing milk and milk products were interviewed from all the five districts in Dar es Salaam. The highest percentage of respondent (63.3%) belonged to the 33-43 age group, followed by 20.0% of respondents who were in (22-32) age group, then 10% respondents were in 44-54 age group and 6.7% respondents were above 55 age group. The results of this study indicated that, 66.7% of respondents were male and only 33.3% of them were female. About 37% of respondents attended college/university education while 46.7% attended secondary school education 16.7% of the respondents attended primary school education and there were no respondents who have informal education (Table 3.1).

Table 3.1: Demographic characteristics of the study respondents

Variable	Description	Frequency (N=30)	Percentage %
District	Ilala	6	20
	Kinondoni	6	20
	Temeke	6	20
	Ubungo	6	20
	Kigamboni	6	20
Gender	Male	20	66.7
	Female	10	33.3
Age categories	22-32	6	20.0
	33-43	19	63.3
	44-54	3	10.0
	Above 55	2	6.7
Education level	Informal/not attended	0	0.0
	Primary	5	16.7
	Secondary	14	46.7
	College/university	11	36.7

3.4.2 Knowledge on aflatoxins contamination

Table 3.2 indicated that all 100% of the study respondents in this study knew contaminants in milk, however none (0.0%) of the study respondents knew aflatoxin was among the contaminants in milk, 86.7% of the study respondents knew milk can be contaminated with drugs residues and 13.3% of the study respondents knew pesticide can contaminate milk.

Moreover, results of this study indicated that 83.3% of the study respondent have ever heard of the word ‘‘aflatoxins’’, whereas only 16.7% of the study respondents had never heard of the word aflatoxin. This was similar to the observation reported by Kagera *et al.*, (2019) that 80% of the respondents in Nairobi, Kenya had heard of aflatoxins; Similarly, Udomkun *et al.*, (2018) reported that 85% of farmers in eastern Democratic Republic of Congo were aware of aflatoxins. The results of this study were slightly lower than that reported by Magembe *et al.*, (2016) where 97.2% of the respondents in Kilosa District in Tanzania had heard of aflatoxins. In contrast, the results of this study were higher than the

studies reported by Lee *et al.* (2017) reported that 9.4% of the respondents from six provinces of Vietnam had ever heard of aflatoxins; Ayo *et al.*, (2018) reported that 29% of livestock farmers in Meru district in Tanzania had heard of aflatoxins; Adekoya *et al.*, (2017) reported that 2% of the respondents in Nigeria had heard of aflatoxins; Mir *et al.*, (2020) reported that 42% of dairy farmers in Punjab had heard of aflatoxins; Nakavuma *et al.*, (2020) reported that 47.73% of poultry farmers and feed processors in Uganda were aware of mycotoxin; Fundikira *et al.*, (2021) reported that 3.3% of spice retailers in Dar es salaam region in Tanzania had heard of aflatoxins; Jere *et al.*, (2020) reported that 15% of food handlers in Salima district in Malawi had heard of aflatoxins and Suleiman *et al.*, (2017) reported that over 80% of the maize producers, sellers, and buyers in Tanzania were not aware of the presence of mycotoxins in foods.

The discrepancy in awareness between this particular study and other reported studies may be attributed to several factors including the nature of the study population. For example, in Tanzania in year 2016 aflatoxicosis outbreak in Chemba, Kiteto, Chamwino, Dodoma and Kondoa districts in Dodoma region (Kamala *et al.*, 2018) and in Kenya Probst *et al.* (2007) reported aflatoxicosis events which killed many people might have contributed to the increased awareness on aflatoxins.

Furthermore, none of the respondents (0.0%) in this study knew milk and milk products such as UHT and pasteurized could be contaminated with aflatoxins and all respondents (100%) were unaware on aflatoxins contamination in milk and none (0.0%) of the respondents were aware of association between aflatoxins in feeds with aflatoxins in milk. This was almost similar to the research carried out by Gizachew *et al.*, (2016) who reported that over 90% of dairy farmers in Ethiopia were unaware of aflatoxin contamination of milk. The level of contamination observed in this study was higher than

that reported by Mir *et al.* (2020) in Punjab whereby 84% of dairy farmers were unaware of aflatoxin contamination of milk; Lee *et al.* (2017) reported that 79.71% respondents from six provinces of Vietnam were unaware of aflatoxin contamination in milk; Udomkun *et al.*, (2018) reported that 46% of farmers in eastern Democratic Republic of Congo were not aware of aflatoxins contamination in milk. Lack of awareness on aflatoxin contamination among milk processors in this study could be a major contributing factor to aflatoxin contamination in milk hence a threat to public health.

This study revealed moderate knowledge on ill-health effects to human and animals caused with consumption aflatoxins, where by 33.3% of the respondents were aware of the health effects to human and animals caused with consumption of aflatoxins contaminated food and 66.7% of the respondents were not aware of the health effects. This could be due to inadequate information on ill health effects caused aflatoxins. The observation made in this study similar to those observations by Magembe *et al.*, (2016) in Kilosa district in Tanzania who reported that 66.7% of respondents were not aware of the ill-health hazards caused by mycotoxins. Results on awareness level in this study were slightly higher than those reported by Mboya and Kolanisi (2014) who observed that that 58.5% of the participants in Rungwe district, Tanzania were not aware of health hazards associated with the ingestion of mycotoxins in food. In contrast, the results on awareness levels of this study were higher than the study conducted by Fundikira *et al.* (2021) who reported that none of the spice retailers in Dar es salaam, Tanzania were aware of the health effects of aflatoxins, and Ngoma *et al.* (2017) reported that only 1.4% of the parents in central part of Tanzania were aware of ill-health effects caused by aflatoxins contamination in complementary foods.

Table 3.2: Knowledge on aflatoxins by the study respondents

Variable	Description	Frequency (N=30)	Percentage %
Knowledge of contaminants in milk	Yes	30	100.0
	No	0	0.0
Contaminants in milk	Drug residue	26	86.7
	Pesticides	4	13.3
	Aflatoxin	0	0.0
Heard of aflatoxin	Yes	25	83.3
	No	5	16.7
Aware of aflatoxin contamination in milk and milk products such as UHT and pasteurized	Yes	0	0.0
	No	30	100.0
Factors for aflatoxin in milk	Cattle consume toxins in feeds	0	0.0
	Do not know	30	100.0
Aware of consumption of contaminated food cause health effect to human and animals	No	20	66.7
	Yes	10	33.3

3.4.3 Milk handling practices

About 73.3% of the milk processors procured raw milk for processing from their own dairy farms and milk collectors, while 26.7% of the processors procured raw milk for processing from their own dairy farms (Table 3.3). Basing on production capacity of milk processors, 46.7% produced less than 100 l/day; 46.7% produced 100-1000 l/day; and (6.7%) produced more than 1000 l/day. It was observed that all milk processors (100%) assessed the raw milk quality parameters at reception before processing; 36.7% assessed colour, 3.3% assessed smell and 60% assessed density. However, none (0.0%) of the respondents analysed raw samples for presence of aflatoxins. This observation might be due to lack of knowledge on the carry-over of aflatoxin from feeds to animal product such as milk among milk processors. This is supported by the fact that most of the aflatoxin's

awareness in Tanzania were conducted in few products such as maize, groundnuts and animal feeds (Ayo *et al.*, 2018; Magembe *et al.*, 2016).

Table 3. 3: Handling practices of milk by the study respondents

Variable	Description	Frequency N=30	Percentage%
Where raw milk is obtained	Own dairy farm	8	26.7
	Dairy farm and milk collectors	22	73.3
Production capacity (Liters) per day	Less than 100	14	46.7
	100-1000	14	46.7
	Greater than 1000	2	6.7
Quality inspection of milk before processing	Yes	30	100.0
	No	0	0.0
Parameters for inspection	Colour	11	36.7
	Smell	1	3.3
	Density	18	60.0
Checking aflatoxin contamination in raw milk before processing	Yes	0	0.0
	No	30	100.0

3.4.4 Feeding practices of lactating cow by the study respondents

All respondents (100%) in this study raised lactating cattle as a source of milk for processing; 50% used hay, 46.7% used cereal products while 3.3% mixed feeds with other products such as wastes from the kitchen for feeding lactating cattle (Table 3.4). The results indicated that all respondents (100%) mixed lactating cattle feeds with concentrates. It was observed that kitchen wastes such as bread and mouldy left overs were used as animal feeds and cereal products like spoilt maize grains were used as cattle feed. This was also observed by Magembe *et al.* (2016) in Kilosa district, Tanzania and Mboya and Kolanisi (2014) in Rungwe district in Tanzania who reported that 18.1% and 38.4% of mouldy maize grains were used as animal feeds respectively. This observation might be a good source of aflatoxins contamination in cattle feeds and milk.

In this study 90% of the respondents did not inspect for mould growth in cattle feeds before feeding the cows and only few (10%) of respondents were carried out physical inspection by smelling, colour and appearance. This is supported by the fact that aflatoxins cannot be inspected visually only by laboratory test. The results of this study were similar to the studies reported by Jere *et al.*, (2020) whereby food handlers in Malawi inspected quality of the grains such as discoloration, damaged and off odour through physical observation and Suleiman *et al.* (2017) in Tanzania reported that quality of the grains was determined by physical observation such as broken grains, infestation and discoloration. Therefore, when these factors considered can reduce level of contamination.

It was observed that all respondents (100%) stored lactating cattle feeds for use during period of scarcity (70%), in times of lower prices (26.7%) and surplus (3.3%). About 77% of the respondents stored cattle feeds for more than one month while 23.3% of the respondents stored cattle feeds for less than one month. It was observed that 23.3% of the study respondents stored cattle feeds in quantity of less than 100 kg; 66.7% of respondents stored cattle feeds ranging from 100 to 1000 kg and 10% of respondents stored cattle feeds in quantity exceeding 1000 kg. Moreover, it was observed that 86.7% of respondents stored lactating cattle feeds on the floor while 13.3% of respondents stock piled cattle feeds. The results indicated that 80.0% of the respondents did not monitor mould growth in cattle feeds during storage while 20% of the respondents monitored mould growth on stored feeds. Only 20% of the respondents dried feeds in order to prevent mould growth while (80%) of the respondents were not aware of any means for prevention/control of mould growth in cattle feeds.

It has been reported that poor storage practices such as stock piling and other poor bulk storage practices of feeds and prolonged time in storage, influenced contamination of feed stuffs with aflatoxigenic producing fungi (Nakavuma *et al.*,2020; Makau *et al.*,2016). Therefore, presence of aflatoxins in feeds may result into aflatoxin M₁ contamination in milk upon lactating cow consumption.

Table 3.4: Feeding practices of lactating cow by the study respondents

Variable	Description	Frequency (N=30)	Percentage %
Have lactating cattle	Yes	30	100.0
	No	0	0.0
Lactating cattle feeds	Hay	15	50
	Cereal products	14	46.7
	Wastes from kitchen	1	3.3
Mixing feeds with concentrates	Yes	30	100.0
	No	0	0.0
Where concentrates purchased	Local supplier	19	63.3
	industries	10	33.3
Inspection of fungal in lactating cattle feeds	Yes	3	10
	No	27	90
Inspection Parameters	Smell	1	3.3
	Colour	2	6.7
	No inspection	27	90.0
Animal feeds storage	Yes	30	100.0
	No	0	0.0
Reason for storage	Period of scarcity	28	70.0
	Lower price	8	26.7
	Surplus	1	3.3
Storage area	Stock piling	4	13.3
	On floor	26	86.7
Storage duration	Less than month	7	23.3
	More than month	23	76.7
Storage quantity (kg)	Less than 100	7	23.3
	Between 100-1000	20	66.7
	Greater than 1000	3	10.0
Monitoring fungal growth	Yes	6	20.0
	No monitoring	24	80.0
Prevention fungal growth	No	24	80.0
	Proper drying	6	20.0

3.4.5 Relationship between knowledge of aflatoxin with age and level of education

Cross tabulation analysis was used to describe and determine the relationship between the two variables. In this study relationship between knowledge of aflatoxins with age and education level of the respondents was investigated.

Table 3.5: Relationship between knowledge on aflatoxins by education level of the study respondents

Education level of the respondents	Have you ever heard about aflatoxins?			<i>P – Value</i>
	Number of respondents	Yes n (%)	No n (%)	
Primary	5	0(0.0)	5 (100)	0.000*
Secondary	14	9 (64.3)	5 (35.7)	
College/University	11	11 (100)	0 (0.0)	
Total	30	20 (66.7)	10 (33.3)	

- Superscript denoted by * indicate significant difference at $P < 0.05$
- n is the number of the study respondents

The relationship between knowledge on aflatoxins with education level of the respondents was investigated (Table 3.5). The results obtained in this study revealed that education level was directly related to aflatoxins awareness and it was statistically significant at $P < 0.05$ whereby respondents who were highly educated had better knowledge of aflatoxins compared to less educated respondents. Respondents who had attended college/university education level were more aware of aflatoxins compared to the respondents who had attended secondary school education, similarly respondents who had attended secondary school education were more aware of aflatoxins compared to the respondents who had attended primary school education. This is due to fact that people who are more educated have more access and tend to seek for more information on food safety and related issues. These findings were similar to studies done by Adekoya *et al.*, (2017); Jere

et al., (2020); Suleiman *et al.*, (2017) and Udomkun *et al.*, (2018) indicating relationship between education level with aflatoxins awareness reported that educated people were more knowledgeable about aflatoxins compared to less educated people.

Table 3.6: Relationship between knowledge of aflatoxins with age of the study respondents

Age of the respondents (years old)	Number of Respondents(n)	Have you ever heard about aflatoxins?		P – Value
		Yes n(%)	No n(%)	
22-32	6	2 (33.3)	4 (66.7)	0.000*
33-43	19	16 (84.2)	3 (15.7)	
44-54	3	1 (33.3)	2 (66.7)	
Above 55	2	1 (50)	1 (50)	
Total	30	20 (66.7)	10(33.3)	

- Superscript denoted by * indicate significant difference at $P < 0.05$
- n is the number of the study respondents

In this study, relationship between knowledge on aflatoxins with age of the respondents was investigated, (Table 3.6) indicated that 33.3% (2/6) of the respondents aged between 22-32 years were aware of aflatoxins; 84.2% (16/19) of the respondents 33-43 years old were aware of aflatoxins; 33.3% (1/3) of the respondents aged between (44-54) years old were aware of aflatoxins and 50.0% (1/2) of the respondents above 55 years old were aware of aflatoxins. The results of this study indicated younger age group had significant high knowledge on aflatoxins contamination at $P < 0.05$ compared to the old age group. This could be due to the fact that younger age groups were more likely to be more educated or informed about aflatoxins. This is supported by the studies done by Ayo *et al.* (2018) and Lee *et al.* (2017) who reported that younger age groups were more aware of aflatoxins and had the ability to access information faster.

3.5 Conclusion

This study gave a highlight into the handling practices of milk, best feeding practices of lactating cow and level of knowledge of milk processors on issues of aflatoxins in milk and milk products. It was observed that, milk processors in the dairy processing had knowledge on aflatoxins and ill-health effects on humans and animals caused by consumption of foods contaminated with aflatoxins. However, none of the respondents had a knowledge of aflatoxin M₁ contamination of milk and milk products such as pasteurized and UHT milk. In addition, this study indicated that there was relationship between knowledge on aflatoxins with education level and age of the respondents.

Furthermore, none of the respondents had knowledge on factors that contribute to aflatoxin M₁ contamination in milk, that lactating cow fed with feeds contaminated with aflatoxins could excrete aflatoxin M₁ in milk. Poor feeding practices and low-quality feeds such as low-quality grains may cause contamination of milk with aflatoxin M₁. This study contributed to knowledge on aflatoxin M₁ content in milk-to-milk value chain stakeholders in Tanzania.

3.6 Acknowledgements

The authors are thankful to Tanzania Bureau of Standards (TBS) for funding this research. The cooperation given by livestock keepers and the milk processors in Kinondoni, Ilala, Temeke, Ubungo and Kigamboni districts, who participated in this study is highly acknowledged.

References

- Achaglinkame, M. A., Opoku, N., and Amagloh, F. K. (2017). Aflatoxin contamination in cereals and legumes to reconsider usage as complementary food ingredients for Ghanaian infants: A review. *Journal of Nutrition and Intermediary Metabolism* 10: 1-7.
- Alshannaq, A., and Yu, J. H. (2017). Occurrence, toxicity, and analysis of major mycotoxins in food. *International Journal of Environmental Research and Public Health* 14(6): 1-20.
- Ayo, E. M., Matemu, A., Laswai, G. H., and Kimanya, M. E. (2018). Socioeconomic characteristics influencing level of awareness of aflatoxin contamination of feeds among livestock farmers in Meru district of Tanzania. *Scientifica* 2018: 1-11.
- Ayofemi Olalekan Adeyeye, S. (2020). Aflatoxigenic fungi and mycotoxins in food: a review. *Critical Reviews in Food Science and Nutrition* 60(5): 709-721.
- Chilaka, C. A., De Kock, S., Phoku, J. Z., Mwanza, M., Egbuta, M. A., and Dutton, M. F. (2012). Fungal and mycotoxin contamination of South African commercial maize. *Journal of Food, Agriculture and Environment* 10(2): 296-303.
- Choudhary, A. K., Tudu, S., Kumari, P., and Ranjan, A. (2020). Present status, prevalence and seasonal variations of aflatoxin in cattle feed, Bihar, India. *Indian Journal of Science and Technology* 13(17): 1738-1745.

- Eskola, M., Kos, G., Elliott, C. T., Hajšlová, J., Mayar, S., and Krska, R. (2020). Worldwide contamination of food-crops with mycotoxins: Validity of the widely cited 'FAO estimate' of 25%. *Critical Reviews in Food Science and Nutrition* 60 (16): 2773-2789.
- Fundikira, S. S., De Saeger, S., Kimanya, M. E., and Mugula, J. K. (2021). Awareness, handling and storage factors associated with aflatoxin contamination in spices marketed in Dar es Salaam, Tanzania. *World Mycotoxin Journal* 14(2): 191-200.
- Gizachew, D., Szonyi, B., Tegegne, A., Hanson, J., and Grace, D. (2016). Aflatoxin contamination of milk and dairy feeds in the Greater Addis Ababa milk shed, Ethiopia. *Food Control* 59: 773-779.
- Gurbuz, Y., Hussein, A., and Ozkose, E. (2019). Determination of AFLD and AFLQ Genes Responsible for Aflatoxin Formation in Some livestock Concentrated Feeds. *Pakistan Veterinary Journal* 39(2): 163-168.
- Haque, M. A., Wang, Y., Shen, Z., Li, X., Saleemi, M. K., and He, C. (2020). Mycotoxin contamination and control strategy in human, domestic animal and poultry: A review. *Microbial Pathogenesis*, 142: 1-12.
- Horky, P., Skalickova, S., Baholet, D., and Skladanka, J. (2018). Nanoparticles as a Solution for Eliminating the Risk of Mycotoxins. *Nanomaterials* 8(9): 1-21.

- IARC Working Group on the Evaluation of Carcinogenic Risks to Humans. (2012). Chemical agents and related occupations. *IARC monographs on the evaluation of carcinogenic risks to humans*, 100(Pt F): 9-562.
- Ismail, A., Gonçalves, B. L., de Neeff, D. V., Ponzilacqua, B., Coppa, C. F., Hintzsche, H., ... and Oliveira, C. A. (2018). Aflatoxin in foodstuffs: Occurrence and recent advances in decontamination. *Food Research International* 113: 74-85.
- Jere, G. M., Abong, G. O., Njue, L. G., Masamba, K., and Omayio, D. G. (2020). Postharvest Handling Knowledge and Practices among Food Handlers on Mycotoxigenic Molds Contamination in Maize Based Diets in School Meals Program in Salima District, Malawi. *Asian Food Science Journal* 16(3): 1-17.
- Kagera, I., Kahenya, P., Mutua, F., Anyango, G., Kyallo, F., Grace, D., and Lindahl, J. (2019). Status of aflatoxin contamination in cow milk produced in smallholder dairy farms in urban and peri-urban areas of Nairobi County: a case study of Kasarani sub county, Kenya. *Infection Ecology and Epidemiology* 9(1): 1-7.
- Kamala, A., Kimanya, M., Haesaert, G., Tiisekwa, B., Madege, R., Degraeve, S., Cyprian, C. and De Meulenaer, B. (2016). Local post-harvest practices associated with aflatoxin and fumonisin contamination of maize in three agro-ecological zones of Tanzania. *Food Additives and Contaminants: Part A*, 33(3): 551-559.
- Kamala, A., Shirima, C., Jani, B., Bakari, M., Sillo, H., Rusibamayila, N., ... and Simba, A. (2018). Outbreak of an acute aflatoxicosis in Tanzania during 2016. *World Mycotoxin Journal* 11(3): 311-320.

- Ketney, O., Santini, A., and Oancea, S. (2017). Recent aflatoxin survey data in milk and milk products: A review. *International Journal of Dairy Technology* 70(3): 320-331.
- Li, S., Min, L., Wang, P., Zhang, Y., Zheng, N., and Wang, J. (2017). Occurrence of aflatoxin M₁ in pasteurized and UHT milks in China in 2014–2015. *Food Control* 78: 94-99.
- Lee, H. J., and Ryu, D. (2017). Worldwide occurrence of mycotoxins in cereals and cereal-derived food products: public health perspectives of their co-occurrence. *Journal of Agricultural and Food Chemistry* 65(33): 7034-7051.
- Lee, H. S., Nguyen-Viet, H., Lindahl, J., Thanh, H. M., Khanh, T. N., Hien, L. T. T., and Grace, D. (2017). A survey of aflatoxin B₁ in maize and awareness of aflatoxins in Vietnam. *World Mycotoxin Journal* 10(2): 195-202.
- Magembe, K. S., Mwatawala, M. W., Mamiro, D. P., and Chingonikaya, E. E. (2016). Assessment of awareness of mycotoxins infections in stored maize (*Zea mays* L.) and groundnut (*arachis hypogea* L.) in Kilosa District, Tanzania. *International Journal of Food Contamination* 3(1): 1-8.
- Mahmoodi Maymand, M., Mazaheri, M., and Talebi Mehrdar, M. (2019). Determination of Aflatoxin M₁ in Pasteurized Liquid and Powdered Milk Products Imported to Iran. *Iranian Journal of Toxicology* 13(2): 19-23.

- Makau, C. M., Matofari, J. W., Muliro, P. S., and Bebe, B. O. (2016). Aflatoxin B₁ and Deoxynivalenol contamination of dairy feeds and presence of Aflatoxin M₁ contamination in milk from smallholder dairy systems in Nakuru, Kenya. *International Journal of Food Contamination* 3(1): 1-10.
- Mboya, R. M., and Kolanisi, U. (2014). Subsistence farmers' mycotoxin contamination awareness in the SADC region: implications on Millennium Development Goal 1, 4 and 6. *Journal of Human Ecology* 46(1): 21-31.
- Mir, I., Singh, P., Bedi, J. S., Kansal, S. K., and Verma, H. K. (2020). Awareness of Aflatoxin Contamination in Feed by Dairy Farmers of Punjab. *International Journal of Current Microbiology and Applied Sciences* 9(1): 2344-2351.
- Mmongoyo, J. A., Wu, F., Linz, J. E., Nair, M. G., Mugula, J. K., Tempelman, R. J., and Strasburg, G. M. (2017). Aflatoxin levels in sunflower seeds and cakes collected from micro-and small-scale sunflower oil processors in Tanzania. *PLoS One* 12(4): e0175801.
- Nakavuma, J. L., Kirabo, A., Bogere, P., Nabulime, M. M., Kaaya, A. N., and Gnonlonfin, B. (2020). Awareness of mycotoxins and occurrence of aflatoxins in poultry feeds and feed ingredients in selected regions of Uganda. *International Journal of Food Contamination* 7: 1-10.
- Ngoma, S. J., Kimanya, M., Tiisekwa, B., and Mwaseba, D. (2017). Perception and attitude of parents towards aflatoxins contamination in complementary foods and its management in central Tanzania. *The Journal of Middle East and North Africa Sciences* 10(4086): 1-16.

- Palumbo, R., Crisci, A., Venâncio, A., Cortiñas Abrahantes, J., Dorne, J. L., Battilani, P., and Toscano, P. (2020). Occurrence and co-occurrence of mycotoxins in cereal-based feed and food. *Microorganisms* 8(1): 1-17
- Probst, C., Njapau, H., and Cotty, P. J. (2007). Outbreak of an acute aflatoxicosis in Kenya in 2004: identification of the causal agent. *Applied and Environmental Microbiology* 73(8): 2762-2764.
- Suleiman, R. A., Rosentrater, K. A., and Chove, B. (2017). Understanding postharvest practices, knowledge, and actual mycotoxin levels in maize in three agro-ecological zones in Tanzania. *Journal of Stored Products and Postharvest Research* 8(7): 73-84.
- Udomkun, P., Wossen, T., Nabahungu, N. L., Mutegi, C., Vanlauwe, B., and Bandyopadhyay, R. (2018). Incidence and farmers' knowledge of aflatoxin contamination and control in Eastern Democratic Republic of Congo. *Food Science and Nutrition* 6(6): 1607-1620.
- Vaz, A., Cabral Silva, A. C., Rodrigues, P., and Venâncio, A. (2020). Detection methods for aflatoxin M₁ in dairy products. *Microorganisms* 8(2): 1-16.
- WHO, (2018): Aflatoxins – Food Safety Digest. [[www.who.int/foodsafety/FSDigest Aflatoxins EN.pdf](http://www.who.int/foodsafety/FSDigest/Aflatoxins%20EN.pdf)] site visited on 18/5/2021.

CHAPTER FOUR

4.0 OVERALL CONCLUSION AND RECOMMENDATIONS

4.1 Conclusions

In this study, the level of AFM₁ in marketed pasteurized and UHT milk in Dar es salaam, Tanzania was found to be rather high and exceeding maximum level of 0.05 µg/L as per EU regulations. However, none of the contaminated sample exceeded Codex limit of 0.5 µg/L, though such contamination does not guarantee the safety of the products to the Tanzania population. Therefore, measures should be taken in order to minimize the level of AFM₁ contamination. Furthermore, samples originating from hot humid coastal climatic zones had high level of contamination of AFM₁ compared to temperate highland zones this is because hot humid coastal zones had favorable conditions for aflatoxigenic fungi to grow, therefore more effort should be taken to these areas to minimize AFM₁ contamination.

Furthermore, lack of knowledge on AFM₁ contamination in milk, poor storage practices of animal feeds and lack of knowledge on health effects caused by aflatoxins among milk processors observed in this study might be a major reason for AFM₁ contamination in milk. Therefore, the best way to deal with this problem is to reduce AFB₁ contamination in animal feedstuffs by improved processing and storage practices. At the same time, attention should be given to regular monitoring of aflatoxins in animal feed and dairy products. In addition, the governmental agencies should train the farmers, dairy companies and dairy product consumers on the potential health consequences of aflatoxins. Finally, milk and dairy products with high levels of AFM₁ must not be allowed for human consumption by the public health authorities.

4.2 Recommendations

Based on the findings of this study the followings are recommendations:

1. In order to preserve the quality and ensure the safety of milk and dairy products consumed in Tanzania, governmental authorities should set a clear strategy that aims at reducing AFM₁ contamination.
2. Regulatory authorities in Tanzania should set limits for AFM₁ in milk and milk products, since both Tanzanian standards and East African Standards not cover recommended level for AFM₁. Therefore, the findings of this study will provide a base in standard development.
3. More studies should be conducted in Tanzania to determine contamination levels of AFM₁ on milk and milk products such as cheese, yogurt, ice cream.
4. Studies to determine levels of aflatoxin B₁ in the animal feeds given to lactating cows should be conducted. Such studies will be helpful to control the health risk factors and supply of AFM₁ free milk to Tanzania population.
5. It is recommended that all stakeholders involved in milk processing should be extensively trained on aflatoxins and proper feeding quality of cattle feeds to prevent AFM₁ contamination in milk.
6. Knowledge on AFM₁ contamination in milk should be extended to all consumers as milk is a food product that is consumed by all age groups. Therefore, all population should be sensitized on aflatoxins ill health effects.

7. Regulatory authorities should reinforce extensive training to dairy farmers and livestock keepers on best practices such as best feeding practices and good storage practices of animal feeds. Sensitization of dairy farmers, milk processors and feed millers could be achieved through regulatory authorities and government agencies.

8. Intervention strategies by regulatory agencies such as Tanzania Dairy Board (TDB) through their enforcement compign should be conducted to all milk stakeholders on knowledge of aflatoxin M₁ contamination in milk and milk products and the associated health effects to human and animals.

APPENDICES

Appendix 1: Photos showing poor storage practices of cattle feeds for feeding cow



Husks and spoilt maize stored on floor were used for feeding cattle



Products from kitchen were used for feeding cattle

Appendix 2: Questionnaire to assess knowledge and practices on aflatoxin contamination in dairy products to milk processors

This questionnaire is aimed to assess milk processor's knowledge on factors and practices associated with aflatoxin contamination in milk. It will take less than thirty minutes to complete this questionnaire. Please note that your answer will be considered completely confidential and your name will not be included in any reports about these results. Also, your answer as individual will not be shared to anyone.

PART A: GEOGRAPHICAL LOCATION

1. Region.....
2. District.....
3. Division.....
4. Ward.....
5. Village.....

PART B: RESPONDENT DETAILS

6. Date of interview: Day.....Month.....Year.....
7. Sex of respondent: Male Female
8. Age of the respondent (years).....
9. Level of education of the respondent
 - Primary education level
 - Secondary school education level
 - Collage/University education
 - Informal/Not attended any school
10. What type of milk product are you processing?

<input type="checkbox"/> Pasteurized milk	<input type="checkbox"/> Yoghurt
<input type="checkbox"/> UHT milk	<input type="checkbox"/> fresh milk
<input type="checkbox"/> Cultured milk	<input type="checkbox"/> All types of milk product
- Any other (Specify).....

PART C: KNOWLEDGE ON AFLATOXIN

11. Do you know contaminants in milk?

Yes No

12. If yes what types of contaminants in milk do you know?

Aflatoxin drug residues
 Metals pesticides
 Any other (specify)

13. Have you ever heard of aflatoxin (in local language??)?

Yes No

14. If yes, where did you first hear about it?

Radio TV Hospital
 Training Others (Specify).....

15. Do you know aflatoxin can contaminate foods and animal feeds?

Yes No

16. If, yes which type of food products are most contaminated with aflatoxin

Cereal products Spices
 Milk oil seeds
 Do not know Any other (specify)

17. What factors contribute to aflatoxin contamination?

Rodents/Insects/Molds		Do not know	
<input type="checkbox"/>	Poor harvesting/ Storage	<input type="checkbox"/>	Others (Specify)
<input type="checkbox"/>	Moisture		

18. Do you know milk and milk product such as UHT and pasteurized can be contaminated with aflatoxin?

Yes No

19. If, Yes mention factor which cause milk contamination with aflatoxin

When animal consume contaminated feed Do not know
Any other (specify)

20. Do you know consumption of milk contaminated with aflatoxin can cause health effect to animals and human?

Yes No

21. If Yes, mention any health effect of aflatoxin contamination to human?

Cancer stunted /poor growth
Do not now Any other (specify)

29. Do you have lactating cattle do you have?

Yes No

30. What are the main types of feeds are you using for feeding lactating cow?

Hay cereal bran
 kitchen wastes residues from breweries industries
 Seed cake Any other (specify).....

31. Do you mix feeds with other supplements (concentrates)?

Yes No

32. If Yes, where do you obtain supplements (concentrates) for feeding lactating cow?

Local supplier Industries

Commercial processor

Any other (specify)

33. Do you check fungal growth in supplements (concentrates)/feeds before feeding lactating cow?

Yes No

34. If **Yes**, how do you check fungal growth of feeds/supplement?

Smelling Appearance, not moldy
 Colour Any other (specify)

35. During checking, have you encountered any fungal/moulds growth in supplements /feeds?

Yes No

36. If **Yes**, how do you manage the problem?

Disposing Proper drying
Binder Purchase from reliable source
Nothing/proceed with feeding
Any other (specify).....

37. Do you store feeds/supplements?

Yes Yes No

38. If **Yes**, what are reasons for storing feeds/supplements?

Preparation for period of scarcity Favorable/lower prices
Others (specify)

39. How do you store feeds/supplements?

stock piling Raised platform
On floor Any other (specify)

40. What is the storage duration?

Less than 1 month more than 1 month

41. What is the storage quantity (kg)?

Less than 100 Greater than 1000
Between 100–1000

42. During storage, have you encountered any fungal/moulds growth in supplements /feeds?

Yes

No

43. If **Yes**, how do you manage the problem?

Fumigation

Proper drying

Binder

Nothing/proceed with storage

Practice first in first out

Disposing

Purchase from reliable source

Any other (specify).....

Thank you very much.