



Effect of Processing Methods on the Micronutrient Profile, Colour, and Anti-nutritive Components of *Justicia heterocarpa* (mwidu)

Zenorina Aloyce Swai ^{a*}, Frida Albinus Nyamete ^a and Valerian. C. K. Silayo ^a

^a *Department of Food Science and Agro-processing, Sokoine University of Agriculture, Morogoro, Tanzania.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ejns/2025/v17i21637>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://pr.sdiarticle5.com/review-history/130549>

Original Research Article

Received: 28/11/2024
Accepted: 01/02/2025
Published: 05/02/2025

ABSTRACT

Justicia heterocarpa (mwidu) is a popular indigenous leafy vegetable picked wild in rural regions of Morogoro, Tanzania, during the wet seasons. This research examined the impact of processing on micronutrients, total phenols, and anti-nutrient content. The fresh leaves (FL) underwent direct shade drying (UBLDDR), blanched shade drying (BLDR), blanched oven drying (BLDO), fermentation (FFL), and gas and microwave cooking (FLCO5, FLCO10, and FLMCO2). Vitamins, chlorophyll, minerals, phenolic compounds, and anti-nutrients were analyzed. All laboratory experiments adhered to procedures and guidelines. The pH dropped more significantly to <3.5 in a

*Corresponding author: Email: Zenorina.swai@sua.ac.tz;

3% salt-3% sugar brine solution with 1.328 ± 0.006 mg/100 g of lactic acid compared to its counterpart. Blanched leaves dried in 5 days, but unblanched leaves took 15 days. Fermented samples demonstrated a notable reduction in total chlorophyll concentration (0.0964 ± 0.075 mg/g) compared to other processing techniques. The nutritional and anti-nutritional composition of *Justicia heterocarpa* showed significant change ($P < 0.05$) depending on processing methods. The results indicated a significant loss of vitamin C in the fermented and ten-minute cooked samples, at 74.57% and 61.64%, respectively. Cooked FLMCO2 (107.4%, 11.29 ± 0.03 mg/100g) and FLCO5 (86.26%, 10.51 ± 0.02 mg/100g) exhibited more than a two-fold increase in beta-carotene compared to fresh leaves (3.67 mg/100g). In comparison to alternative processing procedures, samples cooked for 10 minutes exhibited significant mineral leaching, whereas unblanched direct shade drying preserved the highest mineral concentration. Fermented samples (532.83 ± 14.91 GAE/100 g) exhibited a 64.19% increase in total Phenolic compounds compared to fresh leaves (190.83 ± 14.91 GAE/100 g). Nonetheless, tannins increased by 68.1% (254.44 ± 7.45 GAE/100g) in the fermented samples. Samples exposed to extended cooking (17.8 ± 3.17 mg/100 g) and fermentation (40.28 ± 3.34 mg/100 g) exhibited the lowest levels of phytates. The oxalate concentration was significantly decreased in the cooked samples. *Justicia heterocarpa* may serve as a sustainable food supply in areas of Tanzania experiencing nutritional deficits.

Keywords: Anti-nutrients; indigenous leafy vegetables; *Justicia heterocarpa*; micronutrients; processing.

1. INTRODUCTION

Micronutrient deficiencies are prevalent in many underdeveloped nations, where populations often depend on staple cereals like maize and rice, resulting in minimal dietary diversification (Heaney et al., 2024; Li et al., 2020; Misci et al., 2021). Micronutrient deficiency, or hidden hunger, arises when the intake and absorption of essential vitamins and minerals, including vitamins A, C, zinc, and iron, are inadequate to support optimal health and development (Jeje et al., 2023). Vitamin A deficiency accounts for approximately 1.7% of deaths among children under five in Africa (Irakoze et al., 2021). In Tanzania, the prevalence of anemia is 59% among children aged 6-59 months (Bukenya et al., 2023; Ngungulu et al., 2024), 37.8% among pregnant women in the Mbeya region (John et al., 2023), and 23% among adolescents in boarding schools in Kilimanjaro (Nicholaus et al., 2020). The prevalence of stunting and underweight in Morogoro is notably elevated as a result of nutritional deficiencies. A 24-hour dietary recall indicates that the prevalence rates of anemia, vitamin A deficiency, and zinc deficiency in children are 42.9%, 29.3%, and 24.9%, respectively (Stuetz et al., 2019). The World Health Organization recommends a daily intake of 400 grams of fruits and vegetables for adults to combat hidden hunger (Irakoze et al., 2023; Stoll et al., 2021; Sarfo et al., 2022). Due to challenging economic conditions, numerous rural regions in Sub-Saharan Africa, including Tanzania lacks the financial capacity to afford

such quantities (Bukenya et al., 2023; Irakoze et al., 2023; Sarfo et al., 2022).

Indigenous leafy vegetables (ILVs) represent a valuable resource for enhancing dietary diversity and ensuring food security at the household level in developing countries (Atuna et al., 2022; Chacha & Laswai, 2020; Imathiu, 2021; Mungofa et al., 2022). Recent studies have identified ILVs as a significant source of nutrition (Imathiu, 2021; Mungofa et al., 2022; Issa-Zacharia et al., 2024), economic potential (Henze et al., 2020), and adaptability to climatic conditions (Imathiu, 2021; Mungofa et al., 2022; Bokelmann et al., 2022). Indigenous leafy vegetables, while beneficial, experience considerable post-harvest losses during the rainy season (Imathiu, 2021; Kiringia, 2019) and are often scarce in the dry season due to their wild growth and reliance on natural rainfall (Misci et al., 2021). Furthermore, ILVs contain anti-nutrients that hinder the body's capacity to digest and absorb essential nutrients (Ngungulu et al., 2024; Mwanri et al., 2018). Phytates and oxalates are capable of chelating metal ions, including iron, zinc, and calcium, resulting in the formation of insoluble complexes (Naseem et al., 2023). Wide research has been conducted on the processing and preservation of indigenous leafy vegetables (ILVs) (Heaney et al., 2024; Ngungulu et al., 2024; Irakoze et al., 2023; Tiisekwa et al., 2024). Additionally, the effects of processing on the reduction of anti-nutrients in specific ILVs have also been examined (Ngungulu et al., 2024; Irakoze et al., 2023; Sangija et al., 2022). *Justicia heterocarpa*,

referred to as "mwidu," is a popular indigenous leafy vegetable among the residents of the Morogoro Region in Tanzania (Sangija et al., 2022). This plant flourishes in rainy seasons and grows abundantly in fields and around family compounds (Sangija et al., 2022). The fresh leaves exhibit a notable micronutrient profile (Gowele et al., 2019). *Justicia heterocarpa* is significantly impacted by drought during arid seasons. Furthermore, the restricted preservation methods impede its accessibility and availability in dry seasons. This study examined the impact of processing methods (blanching, drying, fermenting, and cooking) on the retention of nutrients and the reduction of anti-nutritional components in *Justicia heterocarpa*.

2. MATERIALS AND METHODS

2.1 Samples Collection

Justicia heterocarpa leaves were collected from a farmer's plots at Kiroka ward, Morogoro Rural District, Tanzania. Tender leaves were plucked with a sharp knife from each plot during the early morning hours, packaged in polyethylene bags, and immediately transported to the Food Science and Agro-processing laboratory. The leaves were trimmed, sorted, washed, and placed in a perforated vessel for water drainage. The vegetables were thoroughly mixed to obtain a representative sample.

2.2 Sample Preparation

Fresh leaves were washed, drained, and vacuum-packaged in polyethylene bags, then freeze-dried until analysis. In contrast, to obtain the unblanched room-dried leaves, the leaves were washed, drained, and placed on drying racks at room temperature ($25 \pm 2^\circ\text{C}$) until a crispy texture was achieved, which took 15 days. Dried leaves were vacuum-packaged and freeze-dried before the analysis.

Blanched dried leaves were washed, drained, and secured in net cloth, subsequently immersed in boiling water at 100°C for one minute. They were immediately rinsed in ice water for one minute, followed by drying on racks at room temperature for five days. The blanched leaves were subsequently dried in an oven at 40°C for 12 hours. The moisture content was measured using a moisture meter (Denver, IR35M-0002300V1-GERMANY). After attaining a crispy texture, the leaves were vacuum packaged and then freeze-dried before analysis.

For fermented leaves, previously trimmed and sorted vegetables were washed with distilled water and placed in the perforated tray with a clean paper towel to enhance the removal of excess water. Fermentation was done in submerged brine (1:3 vegetables: water). Two distinct concentrations of salt to sugar ratios were employed (3%:3% and 4%:2%) as previously outlined by Sangija et al. (2022) and Stoll et al. (2021) with certain modifications. Fermenting vessels utilized were 800 ml plastic containers with secure seals. Locally produced crock pots, consisting of polyethylene bags filled with water, were utilized to submerge the vegetables, preventing the leaves from floating to the surface and thereby mitigating mold growth during fermentation. The experimental setup was incubated at 25°C , which is the optimal temperature for the growth of fermenting bacteria. Samples were collected at 0, 12, 24, 36, 48, 60, 72, 84, 96, 108, 120, and 144 hours. The efficacy of fermentation was evaluated by monitoring and recording pH and lactic acid levels in triplicate. Subsequent to pH measurement, the samples were promptly freeze-dried, awaiting laboratory examination of lactic acid using a spectrophotometer (JENWAY 6715 UV/Vis U.K.).

Regarding cooked leaves, the cooking time for *Justicia heterocarpa* leaves varied from 3 to 20 minutes, according to pilot data collected from the study area. 500 grams of leaves were boiled in a pot with three hundred milliliters of water on a gas cooker for 5 and 10 minutes, respectively. Furthermore, 500 grams of the identical vegetables were subjected to microwave cooking for a duration of 2 minutes. Finally, the vegetables were cooled to room temperature and transferred, along with the cooking water, into plastic jars with lids for freeze-drying prior to laboratory analysis.

2.3 Chemical Analysis

2.3.1 Determination of nutritional composition

Beta-carotene: Total beta-carotene was done as per (Kimura & Rodriguez-Amaya (2002) with slight modification. Frozen samples were thawed to room temperature ($25 \pm 2^\circ\text{C}$) and homogenized using a blender. A total of 2.000 g of sample was weighed in duplicate in 15 ml screw-capped glass tubes, followed by the addition of 10 ml of acetone (Sigma-Aldrich, USA). The samples were vortexed for a minute, then incubated in a water bath (SW23GB,

JULABO) set at 85°C for 10 min. After that, 120 µl of 80% (w/v) potassium hydroxide (Sigma-Aldrich, USA) was added to the tubes, vortexed for 1 min, and incubated further at 85°C for 5 min. The tubes were transferred in ice to cool before the addition of 4 ml of deionized water. The tubes were then vortexed for 1 min, and 5 ml of hexane (Sigma-Aldrich, HPLC grade, USA) was added to each tube and further vortexed for 1 min before centrifugation at 3000 rpm for 5 minutes. The upper phase of hexane was pipetted into a separate 25 ml labeled flask. Extraction from the pellet was repeated four times using 4x3x3x3 ml hexane, and the extract was combined in the 25 ml labeled flask. Leaves containing higher concentrations of beta carotene were further extracted by carotenoids. Sometimes, more extraction was necessary. 5 ml of deionized water was added into the extracts, mixed with vortex for 1 minute, and centrifuged for 5 mins at 3000 rpm. The hexane layer in each was recovered into a clean test tube, and the hexane was evaporated under nitrogen in the N-Evaporator machine (Organomation, Model OA-8125) with its water bath set at 40°C to dry. Dried extracts in tubes were reconstituted by adding 10 ml of methanol (HPLC grade Sigma Aldrich) to the tubes, mixing homogeneously for 1 min by a vortex. The samples were transferred in 1 ml to HPLC vials. A volume of 30 µl was injected into the HPLC (SPD-20A SHIMADZU CORPORATION, Japan) with the type and conditions explained above. The method's limit of detection and limits of quantification (LOD and LOQ) were 3.1 µg/ml and 5.2 µg/ml, while the percentage recovery was 103%, respectively.

Vitamin C (ascorbic acid): The quantification of *Justicia heterocarpa* ascorbic acid was conducted utilizing the 2,4-dinitrophenylhydrazine (2,4-DNPH) colorimetric assay method as previously described by Khasanah & Riyanto (2023). Each sample was finely chopped, blended, and homogenized using an electric blender. A total of 50 mL of a 3% metaphosphoric acid and 8% acetic acid solution was mixed with 10 mL of the blended sample. The resulting mixture was then diluted to the desired concentration by adding the same metaphosphoric-acetic acid solution to a 100 mL volumetric flask. After filtration, the clear filtrate was collected for vitamin C determination. A few drops of bromine water were added to the filtered sample solution to Oxidize the ascorbic acid to dehydroascorbic acid. Excess bromine was neutralized with 10% thiourea, yielding a clear solution. Standard ascorbic acid solutions were

prepared by diluting the 500 ppm stock to final concentrations of 5, 10, 15, 20, and 25 ppm. For both the oxidized sample and standard solutions, 1 mL of 2,4-DNPH solution was added. These were incubated in a 37°C water bath for 3 hours. After incubation, the solutions were cooled in an ice bath for 30 minutes and then treated with 5 mL of 85% H₂SO₄, resulting in a colored solution. The absorbance was measured at 530 nm, and ascorbic acid content was quantified using a standard calibration curve.

Minerals: The mineral content (Fe, Zn, Ca, and Mg) was analyzed through microwave-assisted acid digestion and ICP-OES, as outlined by Jachimowicz et al. (2021) with some modifications. Approximately 0.5 g of the pulverized plant material sample was weighed into a digestion flask. A 10 mL mixture of concentrated nitric acid (HNO₃) and hydrogen peroxide (H₂O₂) (typically in a 3:1 ratio) was added to the flask, which was then allowed to stand for 4 hours in the fume hood. The flask was placed on a microwave digestion system, and the mixture was heated in a hot block (Hot Block 150 models: SC-154-240) at 95°C and boiled for 30 minutes until it reached 85°C and became apparent, which took 1-2 hours. After digestion, the flask was allowed to cool, and the solution was diluted to a known volume of 50 mL with distilled water in a volumetric flask following filtration. The machine was turned on and allowed to warm up. The digested samples in the test tubes were introduced into the Inductively Coupled Plasma Optical Emission Spectroscopy machine (ICP-OES-5900 Agilent) with the model number (Agilent 5900 SVDV ICP-OES), serial number MY2215CP04, software version 7.6.0.12121, and firmware version 5590. The emission signals from the samples were monitored, and the instrument parameters (e.g., plasma conditions and nebulization rate) were optimized for the analysis. The instrument outputted the concentration of the minerals, which were displayed on the computer.

Determination of total phenolic compounds: The total phenol for the *Justicia heterocarpa* sample was determined by using the Folin-Ciocalteu method as described by Sánchez-Rangel et al. (2013) with some modifications. 80 mg of the sample was added to 2 mL of 50% methanol and extracted by maceration for 2 hours. The resulting mixture was centrifuged at 10,000 rpm for 10 minutes to separate the supernatant, and the volume was adjusted to 5 mL with 50% methanol in a volumetric flask. A 1

mL aliquot of the sample extract was transferred into a test tube, followed by the addition of 9 mL of distilled water and 1.5 mL of Folin-Ciocalteu's reagent. The mixture was incubated at room temperature for 5 minutes. Afterwards, 4 mL of 35% (w/w) Na_2CO_3 was added, and the volume was adjusted to 25 mL with distilled water. The mixture was agitated and left to stand for 30 minutes at room temperature. The absorbance of the sample was measured at 765 nm using a UV-Vis spectrophotometer. Gallic acid was used as a standard for the calibration curve. Total phenolic content was expressed as mg gallic acid equivalents per gram of sample (mg/g).

2.3.2 Determination of anti-nutrient composition

Tannins: The tannin content in each sample was determined using insoluble polyvinyl-pyrrolidone (PVPP), as described by Mohankumar et al. (2018). Specifically, 1.0 mL of the extract dissolved in methanol (1%) was mixed with 100 mg of PVPP. The mixture was vortexed and then incubated for 15 minutes at 4°C. The sample was centrifuged for 10 minutes at 3,000 rpm to separate the precipitated tannins. The clear supernatant obtained after centrifugation contained non-tannin phenolic, which was subsequently analyzed using the Folin-Ciocalteu method to determine total phenolic content. The difference in absorbance values before and after the removal of tannins was regarded as the absorbance value for total tannins in the sample.

Phytates: The phytate content was assessed through the iron-precipitation colorimetric method utilizing thiocyanate detection, following the approach outlined by Mwanri et al. (2018) with certain modifications. Each powdered sample, weighing approximately 0.2 g, was carefully measured into a 125 ml Erlenmeyer flask. To extract the phytic acid, 50 ml of 3% trichloroacetic acid (TCA) was added, and the mixture was swirled by hand for 45 minutes with occasional swirling during the first 30 minutes. After centrifugation, 10 ml of the supernatant was transferred to a 50 ml conical flask. Next, 4 ml of FeCl_3 solution was rapidly added to the aliquot, and the content was heated in a boiling water bath for 45 minutes. After 30 minutes of heating, two drops of 3% sodium sulfate were added to the extract, and the heating process continued. The resulting supernatant was centrifuged for 15 minutes and carefully decanted. The precipitate was washed twice with 20 to 25 ml of 3% TCA by thoroughly dispensing, followed by heating in a

boiling water bath for 10 minutes and centrifugation. This washing process was repeated with water. The precipitate was then dispersed in 27 ml of water and mixed with 3 ml of 1.5 N NaOH. The volume was adjusted to approximately 30 ml with water and heated in a boiling water bath for 30 minutes. The precipitate was filtered through moderately retentive filter paper, such as Whatman No. 2. The precipitate was washed with 70 ml of hot water, and the filtrate was discarded. The precipitate was then dissolved from the filter paper using 40 ml of 3.2 N HNO_3 into a 100 ml volumetric flask. The filter paper was thoroughly washed with multiple portions of water, and the washings were collected in the same flask without exceeding the 100 ml volume. The flask was allowed to cool to room temperature and diluted to volume with water. A 5 ml aliquot was transferred to a 100 ml volumetric flask and diluted to approximately 70 ml. Subsequently, 20 ml of 1.5 M KSCN was added, and the solution was diluted to volume. The resulting color was measured immediately within 1 minute at 480 nm using a spectrophotometer. A reagent blank was included for each set of samples to ensure accuracy. The standard stock solution was prepared by taking 1000 mg/L of phytate (phytic acid) from potassium phytate ($\text{K}_3\text{C}_6\text{H}_6\text{O}_{18}\text{P}_3$). Precisely, 2.064589 g of potassium phytate was weighed and dissolved in a 1000 mL volumetric flask to prepare a working solution of different concentrations. The standard solutions were then subjected to coloration, and the absorbance was measured at 480 nm. A standard plot of concentration against standard absorbance was created, and a linear regression equation was obtained. The concentration of phytates was calculated using the Beer-Lambert equation.

Oxalates: The oxalic acid content was quantified utilizing the UV-Vis spectrophotometric approach, employing iron Ferron reagent detection, as outlined by Salgado et al. (2023), with modifications. A 1.00 g sample of dried ground compost was extracted in a 100 ml conical flask with 10 ml of water and 10 ml of citric acid reagent. The flask was heated on a hot plate, allowing it to simmer for 30 minutes with a small glass funnel placed in the neck of the flask as a condenser. The mixture was filtered through No. 30 Whatman filter paper and thoroughly washed with cold water. The flask was drained, and the filter paper and the precipitate were partially dried at 100°C for one hour before both were returned to the flask. Based on the expected calcium oxalate concentration, 25 ml or

50 ml of 0.4 N hydrochloric acid was added using a pipette (25 ml if the expected concentration was below 1% and 50 ml between 1% and 3%). The solution was heated to near the boiling point and maintained at this temperature for 5 minutes with occasional swirling. The solution was then filtered through a small, dry filter paper. The initial filtrate was discarded, and 4 ml of the remaining filtrate was pipetted into a dry beaker. To the filtrate, 5 ml of iron ferron reagent was added. The solution was protected from direct sunlight, and the absorbance was measured at 525 nm using a UV-Vis spectrophotometer (JENWAY 6715 UV/Vis U.K.) with the instrument zeroed against a reagent blank containing 4 ml of 0.4 N hydrochloric acid and 5 ml of iron Ferron reagent. A standard curve was constructed using solutions with calcium oxalate ($\text{CaC}_2\text{O}_4 \cdot \text{H}_2\text{O}$) concentrations ranging from 0 to 4 mg, dissolved in 4 ml of 0.4 N hydrochloric acid.

2.3.3 Determination of physico-chemical parameters

Chlorophyll content: The chlorophyll content was assessed using acetone extraction and UV-Vis spectrophotometry, following the methodology outlined by Sangija et al. (2022), with modifications. Use a mortar and pestle to crush 1.0 g of the material into a fine pulp with 10 ml of 80% acetone (LOBA, India). The pulp was centrifuged at 4000 rpm at room temperature for 5 minutes, then the green supernatant was transferred to a 50 ml volumetric flask for chlorophyll analysis. The centrifuge tube sediment was scraped and pulverized with the same mortar and pestle and a little 80% acetone to remove the remaining chlorophyll. The supernatant was mixed with the previous supernatant in the volumetric flask after 5 minutes of centrifugation at 4000 rpm at room temperature. Re-extraction was repeated until the residue was green-free. A 50 ml volumetric flask with 80% acetone received the supernatant. The extract was chilled for 10 minutes. With 80% acetone as the blank, a spectrophotometer (JENWAY 6715 UV/Vis U.K.) assessed the extract's absorbance at 663 and 645 nm. Chlorophyll was calculated using the empirical formula;

Chlorophyll a, mg/g tissue = $12.7 (A_{663}) - 2.69 (A_{645}) \times V / 100 \times W$.

Chlorophyll b, mg/g = $22.9 (A_{645}) - 4.68(A_{663}) \times V/100XW$

Total Chlorophyll = Chlorophyll a + Chlorophyll b
A represents absorbance at designated

wavelengths; V is the final volume of the chlorophyll extract; W indicates the fresh weight of the extracted tissue.

Lactic acid: Colorimetric Method for Lactic Acid Quantification Using Ferric Chloride was used as outlined by Borshchevskaya et al. (2016), incorporating specific modifications. Five grams of dried leaf sample were placed in a beaker, to which boiled distilled water was added at a ratio of approximately 1:20 to 1:50, and allowed to cool. Following heating, the extract was allowed to cool to room temperature. The extract was then filtered through filter paper (Whatman No. 1) to remove the solid leaf residue, the filtrate test solution (50 μL) containing lactic acid and 2 mL of a 0.1% iron (III) chloride solution in a cuvette and stirred was conducted at $25 \pm 5^\circ\text{C}$. The reaction was allowed to proceed at room temperature (26.2°C) for 10–15 minutes to ensure the lactic acid fully reacted with the ferric ions, forming a yellow-coloured complex. The absorbance was measured at 390 nm. To create the calibration curve, the absorbance of the lactic acid (1.2 g) with a known concentration of 89% (density = 1.2 g/mL) was measured; a stock solution containing 89 g/L of lactic acid was prepared. From this stock solution, a series of lactic acid solutions with different concentrations was prepared by dilution. A plot of absorbance against concentration was generated. The absorbance of the samples was then measured, and the lactic acid concentration was calculated using the calibration curve based on the Beer-Lambert law.

2.4 Statistical Analysis

All data were subjected to Statistical Package for Social Sciences (SPSS) version 27. A one-way analysis of variance (ANOVA) test was performed following a post-hoc Turkey test, with significant differences determined at a 5% level ($P < 0.05$). All results were expressed as mean \pm SD of triplicate values except for beta carotene and minerals, where the values were duplicate.

3. RESULTS AND DISCUSSION

3.1 Effect of Processing on Physicochemical Properties of *Justicia heterocarpa*

3.1.1 Dynamic trend in pH and lactic acid during spontaneous fermentation

The results of the dynamic trend in pH and lactic acid of *Justicia heterocarpa* leaves are presented

in Table 1. Findings suggest significant trends at $P < 0.05$ for pH and lactic acid during the spontaneous fermentation of *Justicia heterocarpa* leaves exposed to two distinct doses of salt-sugar brine solution. The pH drops from neutral to acidic, measuring 3.89 and 3.52 4% salt-2% sugar and 3% salt-3% sugar, respectively, within 48 hours, with 3% salt-3% sugar brine concentration showed a more significant pH reduction than its counterpart. The variation could be due to the increased sugar content, which promoted bacterial growth by giving fermentation additional substrate. The quick pH decrease establishes acidic environment that suppresses spoiling organisms which cannot tolerate low pH and promotes beneficial bacteria (Irakoze et al., 2023). The study on the fermentation of pumpkin leaves in Kenya reported the spontaneous fermenting microbes present in pumpkin leaves when treated with 3% salt-3% sugar brine solution were *Leuconostoc citreum* at time zero, after 48 hours *Weissella cibaria* prevail the fermentation process, and *Lactiplantibacillus plantarum* finalize the fermentation process (Misci et al., 2021). The work conducted in Tanzania by (Sangija et al., 2022), demonstrated successful spontaneous fermentation of *Solanum* species using a 4% salt-2% sugar brine solution, resulting in a pH reduction to 3.7 within 48 hours. Acidification of the fermented vegetables is very essential for preservation of the vegetable and extend its shelf life also achieving the desirable flavor (Degrain et al., 2020).

In comparison to a 4% salt-2% sugar brine solution, a 3% sugar-3% salt brine solution exhibited a considerable rise in lactic acid content over time during spontaneous fermentation, indicating enhanced efficacy in producing an acceptable quantity of lactic acid. Upon completion of fermentation, a 3% sugar-3% salt brine solution yielded 1.3288 mg/100 ml of lactic acid, which was substantially greater ($P < 0.05$) than the 4% salt-2% sugar brine solution, which produced 0.7877 mg/100 ml (Table 1). The lactic acid produced acts as a preservative of the vegetables because spoilage and pathogenic microbes cannot tolerate high acid. However, it contributes to flavor and aroma development, improving fermented vegetables' sensory appeal (Irakoze et al., 2023). The study conducted by Sangija et al. (2022) on the fermentation of nightshade leaves in 4% and 2% salt sugar brine solution, utilizing the titration method, yielded results comparable to those of the current study.

3.1.2 Moisture content (MC) of dried *Justicia heterocarpa*

The moisture content changes with time for unbranched leaves directly dried in the room (UNBLDR), blanched leaves dried in the room (BLDR), and blanched leaves dried in the oven (BLDO) during the drying of *Justicia heterocarpa* are displayed in Table 2. All three methods displayed substantial moisture reduction of moisture content, in which moisture was reduced from 87.83 to 12.38% for UBLDDR, 85.4 to 10.29 for BLDR, and 86.11 to 9.89%. Samples of blanched leaves dried at room (BLDR) temperature for 48 hours exhibited a significantly higher moisture content ($P < 0.05$) compared to those unbranched leaves dried directly at room temperature (UBLDDR). The study by Traoré et al. (2017) indicated a reduction in moisture content to 11.82% in black nightshades and amaranth when dried in shade. This finding is consistent with the current study, which observed that *J. Heterocarpa* achieved a moisture content of 12.38% under similar drying conditions. While oven drying proved to be the most effective drying method (9.89%, 12 hrs) in the current study due to its efficiency, it is less feasible in resource-limited settings as it necessitates electricity and specialized equipment. Blanching facilitates moisture reduction by disrupting cell walls, thereby enhancing the rate of moisture escape. Achieving the desired moisture content in unbranched vegetables presents greater challenges. The process required 15 days to achieve a moisture content of 12.38%, rendering it less suitable for rapid processing, though viable under resource constraints. Other studies have highlighted the importance of blanching before drying, which leads to minimal nutrient leaching and preserves more nutrients (Fombang & Mbofung, 2017).

3.1.3 Effects of processing on chlorophyll content

Chlorophyll content of fresh and processed leaves of *Justicia heterocarpa* is presented in Table 3. The total chlorophyll content of the Leaves subjected to different processing methods varied and were significantly reduced to 0.0964 ± 0.075 mg/g in fermented samples and 6.7688 ± 0.448 mg/g for the ten-minute cooked leaf samples (Table 3). Total chlorophyll levels were highest in fresh leaves (FL), measuring 13.429 ± 0.522 mg/g. Leaves dried at room temperature (BLDR) and those microwaved for 2 minutes (FLMCO2) maintained total chlorophyll

levels of approximately 12.92 and 12.15 mg/g, respectively, and did not show a significant difference ($p>0.05$) compared to fresh leaves. Leaves dried at room temperature retained a higher chlorophyll level than all other processing methods, which suggests that blanching before drying can control chlorophyll degradation. Fermented fresh leaves (FFL) samples exhibited significantly reduced chlorophyll levels of 0.0964 ± 0.075 mg/g (Table 3), indicating substantial degradation of chlorophyll during acidic fermentation, resulting in the transformation into non-green compounds and observable color changes. Numerous studies indicate a significant decrease in chlorophyll levels in fermented ILVs.

The study conducted by Sangija et al. (2022) on the impact of fermentation on chlorophyll reduction in nightshade species indicated a gradual decrease in chlorophyll a, b, and total chlorophyll levels (Degrain et al., 2020) and reported that the accumulation of acids such as acetic and lactic is the primary factor contributing to the reduction of chlorophyll levels in fermented nightshades. Chlorophyll present in green leafy vegetables serves as a significant antioxidant, aiding in the mitigation of oxidative stress and the prevention of cellular damage. Due to its anti-inflammatory properties, it may reduce chronic inflammation and enhance overall health (Martins et al., 2023).

Table 1. Lactic acid and pH dynamics in spontaneous fermentation (SF)

Time	Lactic Acid (mg/100ml)		P ^H	
	4%salt-2%sugar	3%salt-3%sugar	4%salt-2%sugar	3%salt-3%sugar
0 hrs.	0.099 ± 0.011^a	0.06 ± 0.011^a	6.68 ± 0.006^a	6.62 ± 0.021^a
12 hrs.	0.125 ± 0.006^{ab}	0.12 ± 0.011^b	5.23 ± 0.01^{ab}	5.66 ± 0.023^a
24 hrs.	$0.177 \pm 0.011^{b^*}$	0.22 ± 0.006^c	4.45 ± 0.044^{ab}	4.35 ± 0.031^{ab}
36 hrs.	$0.26 \pm 0.023^{c^*}$	0.35 ± 0.006^d	4.0 ± 0.02^{abc}	3.87 ± 0.02^b
48 hrs.	0.28 ± 0.042^{cd}	0.36 ± 0.011^d	3.89 ± 0.012^{bcd}	3.53 ± 0.023^b
60 hrs.	0.29 ± 0.006^{cd}	0.42 ± 0.040^e	3.82 ± 0.02^{cde}	3.4 ± 0.012^c
72 hrs.	0.33 ± 0.006^{de}	0.55 ± 0.022^f	3.78 ± 0.02^{de}	3.2 ± 0.015^c
84 hrs.	0.35 ± 0.011^e	0.65 ± 0.006^g	3.7 ± 0.0^{ef}	3.31 ± 0.006^c
96 hrs.	0.36 ± 0.017^e	0.75 ± 0.011^h	3.77 ± 0.026^f	3.40 ± 0.015^d
108 hrs.	0.42 ± 0.028^e	0.89 ± 0.006^i	3.68 ± 0.025^g	3.39 ± 0.012^e
120 hrs.	0.52 ± 0.011^f	0.96 ± 0.011^j	3.66 ± 0.057^h	3.27 ± 0.058^f
132 hrs.	0.69 ± 0.017^g	1.20 ± 0.011^k	3.72 ± 0.012^i	3.30 ± 0.006^g
144 hrs.	0.78 ± 0.028^h	1.328 ± 0.006^l	3.64 ± 0.032^j	3.21 ± 0.023^h

Values are means \pm SD and values within the same column with different superscript letters are significantly different from each other at $P<0.05$

Table 2. Moisture content of different drying methods

Time	Moisture%		
	UBLDDR	BLDR	BLDO
0 hours	87.83 ± 1.444 g	84.33 ± 1.624^e	86.11 ± 0.596
12 hours	73.35 ± 2.024^f	43.86 ± 3.879^d	9.89 ± 0.157
24 hours	51.00 ± 2.036^e	28.07 ± 1.356^c	
48 hours	19.08 ± 0.251^d	15.41 ± 1.181^b	
72 hours	16.58 ± 0.592^{cd}	13.06 ± 0.648^{ab}	
96 hours	15.05 ± 0.196^{abc}	11.95 ± 0.205^{ab}	
120 hours	15.15 ± 0.933^{abc}	10.29 ± 0.121^a	
144 hours	15.73 ± 0.529^{bc}		
168 hours	14.21 ± 0.754^{abc}		
Day 8	13.68 ± 0.7^{abc}		
Day 12	14.04 ± 0.072^{abc}		
Day 13	13.44 ± 0.516^{ab}		
Day 14	12.77 ± 0.252^a		
Day 15	12.38 ± 0.27		

Values are means \pm SD and values within the same column with different superscript letters are significantly different from each other at $P<0.05$. Abbreviations: BLDR: Blanched leaves dried in room, BLDO: Blanched leaves dried in oven, UBLDDR: Un-branched leaves directly dried in the room

Table 3. Chlorophyll content in raw and processed *Justicia heterocarpa*

Sample	Chlorophyll a (mg/g)	Chlorophyll b (mg/g)	Total Chlorophyll (mg/g)
FL	8.019 ± 0.104 ^d	5.414 ± 0.422 ^d	13.429 ± 0.522 ^f
BLDR	7.927 ± 0.108 ^d	4.995 ± 0.795 ^{cd}	12.918 ± 0.836 ^{ef}
BLDO	6.403 ± 0.956 ^c	4.316 ± 0.117 ^{cd}	10.715 ± 0.942 ^{cd}
UBLDDR	7.35 ± 0.36624 ^{cd}	4.0759 ± 0.151 ^c	11.4228 ± 0.381 ^{cde}
FFL	0.1065 ± 0.126 ^a	-0.01 ± 0.0530 ^a	0.0964 ± 0.075 ^a
FLCO5	6.3389 ± 0.551 ^c	3.8569 ± 0.790 ^{bc}	10.193 ± 1.066 ^c
FLCO10	3.945 ± 0.5236 ^b	2.8258 ± 0.173 ^b	6.7688 ± 0.448 ^b
FLMCO2	7.7804 ± 0.294 ^d	4.3774 ± 0.235 ^{cd}	12.1545 ± 0.527 ^{def}

Values are means ± SD and values within the same column with different superscript letters are significantly different from each other at $P < 0.05$. Abbreviations: FL: Freshly leaves, BLDR: Blanched leaves dried in room, BLDO: Blanched leaves dried in oven, UBLDDR: Un-branched leaves directly dried in the room, FFL: Fermented fresh leaves; FLCO5: 5 min cooked sample in gas cooker, FLCO10: 10 min cooked sample in gas cooker, FLMCO2: 2 min microwaving cooked samples

Table 4. Beta carotene and ascorbic acid of *Justicia heterocarpa* processed leaves (FW)

	Ascorbic acid (mg/100g)	% Reduction	Beta carotene (mg/100g)	% Reduction
FL	23.21 ± 0.742 ^e	-	3.67 ± 0.072 ^c	-
BLDR	18.62 ± 0.557 ^d	19.79	2.06 ± 0.038 ^b	43.82
BLDO	16.52 ± 0.185 ^{cd}	28.84	5.72 ± 0.156 ^d	-55.79
UBLDDR	19.41 ± 0.186 ^d	16.40	1.55 ± 0.013 ^a	57.78
FFL	5.88 ± 0.743 ^a	74.66	1.29 ± 0.053 ^a	64.69
FLCO5	14.02 ± 1.114 ^c	39.59	10.51 ± 0.002 ^f	-186.26
FLCO10	8.90 ± 1.299 ^b	61.65	9.14 ± 0.035 ^e	-148.77
FLMCO2	18.22 ± 0.371 ^d	21.49	11.29 ± 0.030 ^g	-207.42

Values are means ± SD and values within the same column with different superscript letters are significantly different from each other at $P < 0.05$. % Reduction: The percentage of ascorbic acid and beta carotene. Abbreviations: FL: Freshly leaves, BLDR: Blanched leaves dried in room, BLDO: Blanched leaves dried in oven, UBLDDR: Un-branched leaves directly dried in the room, FFL: Fermented fresh leaves; FLCO5: 5 min cooked sample in gas cooker, FLCO10: 10 min cooked sample in gas cooker, FLMCO2: 2 min microwaving cooked samples

3.2 Effect of Processing on *Justicia heterocarpa* Nutritional Composition

Ascorbic acid: Table 4 presents the beta carotene and ascorbic acid content of processed *Justicia heterocarpa* leaves. Unprocessed fresh leaves (FL) exhibited a high concentration of ascorbic acid (23.21 mg/100 g), which significantly decreased ($P < 0.05$) following various processing methods.

The fermentation of fresh leaves and gas cooking for ten minutes resulted in a significant reduction of ascorbic acid content, approximately by 74.66% and 61.65%, respectively. Various processing methods resulted in a moderate decrease in ascorbic acid levels. Specifically, BLDO decreased ascorbic acid by 28.84%, FLCO5 by 39.59%, and FLMCO2 by 21.49%. In contrast, BLDR and UBLDDR led to reductions of only 19.79% and 16.40%, respectively. Despite the lack of significant difference ($p > 0.05$), the loss of ascorbic acid in blanched dried in oven

leave samples (BLDO) was greater (28.84%) compared to the blanched leaves dried at room temperature (BLDR; 19.79%). Blanching prior to drying enhances the retention of ascorbic acid by inactivating oxidative enzymes, such as ascorbic acid oxidase, responsible for its degradation. Oven drying is a rapid process; however, it results in a significant degradation of ascorbic acid when compared to room drying (BLDR). This degradation may be attributed to elevated thermal temperatures, which enhance the rates of oxidation. The current study achieved a reduction of approximately 75% in ascorbic acid through the fermentation process of *Justicia heterocarpa*. This reduction may be attributed to the solubility of ascorbic acid (vitamin C) in brine solution, as ascorbic acid is soluble in water (Gowele et al., 2019). The findings align with those reported by Sangija et al. (2022) in their study conducted in Tanzania, which indicated a significant loss of vitamin C, ranging from 88.33% to 95%, during the fermentation of *Solanum* species. Furthermore, the gas cooking

of flesh leaves for ten minutes resulted in approximately a 61% reduction in ascorbic acid in this study, likely due to leaching and thermal degradation. A prior study indicated that pasteurizing yellow and red tomatoes at 100°C for 10 minutes resulted in an up to 80% reduction in vitamin C, while cooking broccoli for 5 minutes was associated with a loss of approximately 66% of this vitamin (Fombang & Mbofung, 2017; Mieszczakowska-Fraç et al., 2021). The findings of vitamin C content of *J. Heterocarpa* from the present study are less than those of the daily recommended value for adult men and women (110 mg/day and 95 mg/day, respectively) (Mieszczakowska-Fraç et al., 2021). Traditionally, *Justicia heterocarpa* is cooked as a composite dish mixed with legumes (cowpea, pigeon peas, beans) and okra (author's unpublished surveyed data). This combination can boost the level of vitamin C to meet the required amount in a day. *Justicia heterocarpa* was reported to be similar to African nightshade (*Solanum nigrum complex*) in taste (Sangija et al., 2022) and that incorporating *Solanum species* or *Justicia heterocarpa* into meat, other vegetables, and legumes such as cowpea improves taste, appetite, and increases carotenoids and vitamin C (Oluoch & Habwe, 2012).

Beta-carotene: Beta-carotene levels differed substantially across all processing processes at $P < 0.05$, with the cooked samples FLCO5 and FLCO10 having the greatest quantities (10.5115 mg/100 g and 11.2885 mg/100 g, respectively) (Table 4). The FFL and UBLDD samples exhibited the lowest concentrations of beta carotene, with values of 1.30 mg/100 g and 1.55 mg/100 g, respectively, and were not statistically different from one another. The cooked samples show an increase in beta-carotene content, with FLCO5 exhibiting an 86.26% increase, FLCO10 a 48.77% increase, and FLMCO2 a 107.42% increase. Additionally, the blanched oven-dried vegetables (BLDO) showed a 55% increase in beta-carotene content. The substantial increase in beta-carotene levels in cooked samples may be attributed to the relative heat stability of beta-carotene, as the plant matrix facilitates its release through cell wall breakdown, thereby enhancing its bioavailability (Gowele et al., 2019). This study demonstrated a reduction of approximately 65% in beta-carotene via the fermentation process of *Justicia heterocarpa*. This reduction can be attributed to the sensitivity of beta-carotene to oxidation, which affects its stability over time. Zong et al. (2021) reported

that submerged fermentation of vegetables reduces approximately 20% of total beta-carotene content. However, the study by Sangija et al. (2022) highlighted an increase in beta-carotene during the fermentation of nightshade species. Findings from the present study showed that microwaving cooking resulted in more than a threefold increase in this vitamin. Several studies have reported similar outcomes, with Piyarach et al. (2020) noting that microwaving selected ILVs retained more beta-carotene compared to their fresh counterparts. The observed reduction of beta-carotene in room temperature dried samples (UBLDDR and BLDR) may be attributed to the oxidation of beta-carotene. Research by Piyarach et al. (2020) similarly indicated a reduction in beta-carotene levels in certain vegetables dried at 65°C using a hot air dryer. The study by Ahmed et al. (2023) reported an increase in beta-carotene content in African nightshade and cowpea leaves subjected to various sun-drying treatments.

Minerals: Macro minerals (calcium, magnesium) and trace minerals (iron, zinc) exhibited significant differences across all processing techniques at ($P < 0.05$). Considerable concentrations of Fe (9.46-16.74 mg/100 g), Zn (3.041-4.39 mg/100 g), Ca (255.2-348 mg/100 g), and Mg (71.779-159.9 mg/100 g) were recorded (Table 5).

Iron: Unprocessed fresh leaves (FL) demonstrated a high iron concentration of 16.74 mg/100 g, which significantly decreased ($P < 0.05$) after certain processing methods of *Justicia heterocarpa*. The lowest iron content was observed in samples cooked for ten minutes (FLCO10, 9.46 mg/100 g), indicating significant iron leaching during prolonged cooking. The iron content among the samples (BLDO, BLDR, FFL, and FLCO) exhibited no significant variation, indicating the stability of iron levels across certain processing methods. Irakoze et al., (2023) reported a decrease in iron content in fermented African spider plant and African black nightshade, which aligns with the current study. Conversely, the study by Sangija et al. (2022), utilizing energy-dispersive X-ray fluorescence equipment, reported an increase in iron content in fermented *Solanum species*. This study corroborates findings indicating that iron is susceptible to leaching during cooking and blanching (Kandong, 2019). Iron is essential in the human body for hemoglobin-mediated oxygen transport, energy synthesis, immune function, and muscle performance.

Additionally, it contributes to DNA synthesis and cognitive development (Ogbonnaya et al., 2024). The intake of 100 grams of *Justicia heterocarpa* provides the daily iron requirement for the body, supplying 18 mg for adult women and 6-12 mg for adult men (Stevens et al., 2022).

Zinc: Unprocessed fresh leaves (FL) exhibited a zinc concentration of 4.39 mg/100 g, which significantly decreased ($P < 0.05$) following particular processing methods of *Justicia heterocarpa*, as indicated in Table 5. The study indicated no significant difference in zinc content between blanched dried and fermented samples, likely due to the relative stability of zinc across various processing methods. Furthermore, there is no significant difference between gas-cooked samples subjected to different durations (FLCO5 and FLCO10). A minimal reduction of zinc was observed in microwaved cooked and unblanched room-dried samples, likely due to the absence of liquid, which reduces mineral loss. The zinc content of the fresh leaves in this study is comparable to that reported by Gowele et al. (2019) for the same vegetable (3.9 mg/100 g, FW). The research by Marynurce (2023) indicated a decrease in zinc levels across various drying techniques of nightshade species, consistent with the current study. However, the study by Alassane et al. (2022) reported a contrasting finding, indicating that zinc content increased more than threefold in selected indigenous leafy vegetables (ILVs) dried using sun, oven, and shade methods. Furthermore, the research by Bighaghire et al. (2021) indicated a reduction in zinc content in blanched sun-dried selected indigenous leafy vegetables by as much as 50%. Ukom et al. (2023) reported a decrease in zinc content in the selected cooked indigenous vegetables. Sangija et al. (2022) reported an increase in zinc content in fermented nightshade species, which contradicts the findings of the current study. The zinc content in fresh leaves of *Justicia heterocarpa* exceeds that found in soy (3.7 mg/100 g), indicating a significant mineral concentration in this vegetable (Natesh et al., 2017). Zinc is an essential trace element involved in numerous physiological processes in the body. Its participation in over 300 enzymatic activities and metabolic processes underscores its essential role in maintaining overall health and well-being (Schoofs et al., 2024). Excessive zinc consumption has been associated with health risks, including zinc-induced copper deficiency, symptoms of anemia, and neutropenia (Schoofs et al., 2024).

Calcium: Unprocessed fresh leaves (FL) demonstrated a calcium concentration of 348.73 mg/100 g, which significantly decreased ($P < 0.05$) after specific processing methods of *Justicia heterocarpa*, as shown in Table 5. There was no significant variation ($p < 0.05$) in calcium content observed between shade-dried samples (BLDR and UBLDDR) and high thermal-treated samples (BLDO and FLMCO2). This stability can be attributed to calcium's resistance to changes during various processing methods, as it is not sensitive to heat, light, or oxygen. The research conducted by Alassane et al. (2022) revealed a significant increase in calcium content, exceeding threefold, in selected indigenous leafy vegetables (ILVs) dried through sun, oven, and shade methods. The study by Bighaghire et al. (2021) demonstrated a reduction in calcium content in blanched sun-dried selected indigenous leafy vegetables by up to 50% (Ukom et al. (2023) observed a reduction in calcium content in the chosen cooked indigenous vegetables, consistent with the findings of the current study. Irakoze et al. (2023) reported a decrease in calcium content in fermented African black nightshade and African spider plant, which supports the findings of the current research. Sangija et al. (2022) reported an increase in calcium content in fermented nightshade species, which contradicts the findings of this study. Numerous studies indicate that the calcium content in ILVs is higher than that observed in the current study (Duguma, 2020; Tanor et al., 2020). Calcium is vital for bone and dental health, regulates blood coagulation, influences muscle and nerve function, and activates various enzymes, such as ATPases, crucial for energy production (Tanor et al., 2020). Daily calcium intake should be between 1000 and 1300 mg (Ogbonnaya et al., 2024).

Magnesium: Unprocessed fresh leaves (FL) exhibited a magnesium concentration of 159.9 mg/100 g, which significantly decreased ($p < 0.05$) following specific processing methods of *Justicia heterocarpa*, as indicated in Table 5. No significant variation was observed between samples cooked in a microwave and those cooked with gas for ten minutes (FLMCO2 and FLCO10), as magnesium remains stable and does not degrade or volatilize at standard cooking temperatures. Fermented samples exhibit a notably low magnesium content of 71.78 mg/100 g, likely due to magnesium leaching into the brine solution. Ukom et al. (2023) reported a slight reduction in magnesium content in blanched and sautéed indigenous

Table 5. Mineral content of raw and processed *Justicia heterocarpa* in fresh weight

Sample	Minerals (mg/100g)			
	Iron	Zinc	Calcium	Magnesium
FL	16.74 ± 0.353 ^d	4.39 ± 0.1718 ^d	348.73 ± 0.57 ^f	159.91 ± 0.04 ^g
BLDR	13.40 ± 0.472 ^c	3.58 ± 0.0665 ^b	312.74 ± 0.9 ^d	123.86 ± 0.04 ^b
BLDO	13.84 ± 0.182 ^c	3.4 ± 0.087 ^b	338.30 ± 1.34 ^e	135.30 ± 0.02 ^c
UBLDDR	15.85 ± 0.088 ^d	3.96 ± 0.0411 ^c	314.17 ± 0.1 ^d	149.4 ± 0.66 ^f
FFL	14.25 ± 0.266 ^c	3.53 ± 0.0553 ^b	262.36 ± 0.28 ^b	71.78 ± 0.08 ^a
FLCO5	11.32 ± 0.242 ^b	3.28 ± 0.0661 ^a	286.46 ± 2.12 ^c	140.18 ± 0.023 ^e
FLCO10	9.46 ± 0.354 ^a	3.04 ± 0.0738 ^a	255.20 ± 1.57 ^a	139.09 ± 0.04 ^d
FLMCO2	13.46 ± 0.512 ^c	4.04 ± 0.0019 ^c	339.85 ± 1.51 ^e	138.92 ± 0.06 ^d

Values are means ± SD and values within the same column with different superscript letters are significantly different from each other at $P < 0.05$. Abbreviations: FL: Freshly leaves, BLDR: Blanched leaves dried in room, BLDO: Blanched leaves dried in oven, UBLDDR: Un-branched leaves directly dried in the room, FFL: Fermented freshly leaves; FLCO5: 5 min cooked sample in gas cooker, FLCO10: 10 min cooked sample in gas cooker, FLMCO2: 2 min microwaving cooked samples

Table 6. TPC and anti-nutrient profiles in raw and processed *Justicia heterocarpa*

Sample	Ant-nutrient (FW)			
	Total phenol (GAE/100g)	Tannins (GAE/100g)	Phytates (mg/100g)	Oxalates (mg/100g)
FL	190.83 ± 14.91 ^c	81.17 ± 3.56 ^d	103.56 ± 4.68 ^d	560.46 ± 19.16 ^f
BLDR	142.95 ± 5.92 ^b	59.51 ± 2.96 ^c	76.24 ± 7.83 ^c	351.39 ± 7.41 ^d
BLDO	130.42 ± 5.22 ^b	53.24 ± 2.61 ^c	73.82 ± 5.99 ^c	242.44 ± 4.50 ^c
UBLDDR	352.71 ± 12.01 ^d	164.39 ± 6.01 ^e	94.92 ± 1.80 ^d	417.16 ± 4.50 ^e
FFL	532.83 ± 14.91 ^e	254.44 ± 7.45 ^f	40.28 ± 3.34 ^b	225.76 ± 10.34 ^{bc}
FLCO5	117.19 ± 7.90 ^b	21.32 ± 2.61 ^b	50.66 ± 6.25 ^b	171.77 ± 50.34 ^{ab}
FLCO10	64.75 ± 7.90 ^a	4.22 ± 1.98 ^a	17.81 ± 3.17 ^a	151.16 ± 6.13 ^a
FLMCO2	149.11 ± 29.81 ^b	25.31 ± 7.45 ^b	75.9 ± 1.59 ^c	219.87 ± 1.7 ^{bc}

Values are means ± SD and values within the same column with different superscript letters are significantly different from each other at ($P < 0.05$) Abbreviations: FL: Freshly leaves, BLDR: Blanched leaves dried in room, BLDO: Blanched leaves dried in oven, UBLDDR: Un-branched leaves directly dried in the room, FFL: Fermented fresh leaves; FLCO5: 5 min cooked sample in gas cooker, FLCO10: 10 min cooked sample in gas cooker, FLMCO2: 2 min microwaving cooked samples

vegetables, which is consistent with the findings of the current study. Factors influencing magnesium bioavailability include dietary fat content, the presence of anti-nutrients, and the individual's age and sex (Gowele et al., 2019). Magnesium facilitates energy production, the transport of calcium and potassium ions, nerve conduction, muscular contraction, vasomotor tone, and the maintenance of regular heart rhythm. It serves as a crucial cofactor in over 300 enzymatic systems (Farak et al., 2023). The daily recommended intake of magnesium is between 300 and 429 mg (Farak et al., 2023).

3.3 Effect of Processing on Total Phenolic Compounds (TPC) and Anti-nutrient Composition of *Justicia heterocarpa*

The anti-nutrient profiles of *Justicia heterocarpa* exhibited significant variation under different

processing methods ($P < 0.05$), as indicated in Table 6. The reduction of oxalates, tannins, and phytates following processing improves the bioavailability of minerals, thereby supporting its potential application in addressing micronutrient deficiency.

Total phenolic compounds (TPC): The concentration of Total Phenolic Compounds (TPC) in unprocessed leaves of *Justicia heterocarpa* varied significantly across different processing methods, as indicated in Table 6. No significant variation in TPC was observed in the BLDR, BLDO, and FLCO5 samples. The highest levels of TPC were observed in FFL, which exhibited an increase of 64.19% from the unprocessed leaves, and in UBLDDR, where the increase was 45.9%. The increase in Total Phenolic Compounds (TPC) in unblanched leaves dried at room temperature may be

attributed to the concentration of TPC during the drying process. The increase in TPC in the FFL sample may result from the liberation of bound phenols, enhancing their concentration and bioavailability (Verardo et al., 2020). The low pH stabilizes phenolic compounds, which can be detected in significant quantities (Degrain et al., 2020). Reduced phenol levels following blanching (BLDR) and extended cooking (FLC10) suggest heat sensitivity and potential loss of phenolic components due to oxidation (Mohankumar et al., 2018). Total phenolic compounds exhibit various health benefits, such as antioxidant activity, anti-inflammatory properties, and antimicrobial effects (Leonard et al., 2021).

Tannins: The concentration of total tannins in unprocessed leaves of *Justicia heterocarpa* exhibited significant variation across various processing methods as shown in Table 6. No significant variation was observed between the short time cooked samples (FLCO5 and FLMCO2), likely due to their comparable cooking times, which resulted in similar tannin binding to the food matrix, as well as between the blanched dried samples (BLDR and BLDO). The fermented leaves (FFL) and UBLDDR exhibited the highest tannin concentrations, measuring 254.44 GAE/100g and 164.39 GAE/100g, respectively. An increase of 68.1% in the total tannins in the FFL sample compared to the fresh sample may result from spontaneous fermentation of *Justicia heterocarpa* leaves. During this process, microbial activity and enzymatic actions break down complex tannin compounds and degrade cell walls, thereby releasing bound tannins into a more measurable and soluble form. The ten-minute cooked samples exhibited the lowest tannin levels at 4.22 mg/100 g, reflecting a reduction of 94.6% compared to the two- and five-minute cooking durations. This demonstrates the efficacy of extended cooking in diminishing tannin levels. Irakoze et al., (2023) reported comparable findings, indicating that tannin content was significantly elevated in spontaneously fermented African black nightshade and African spider plants. In contrast to this study, the research conducted by Sangija et al. (2022) indicated a reduction in total tannins in fermented *nightshade species*. A study by Kandonga (2019) reported reduced tannin levels in false sesame and common bean leaves when exposed to varying blanching times and drying techniques, corroborating the findings of the present study. Tannins, due to their pronounced properties and capacity to bind with proteins, can

influence nutrient absorption efficiency. Tannins can interact with proteins via covalent and hydrogen bonding, resulting in the formation of a complex protein precipitate that inhibits absorption (Natesh et al., 2017).

Phytates: The levels of phytates in unprocessed leaves of *Justicia heterocarpa* demonstrated considerable significant variation at ($P<0.05$) among different processing methods as indicated in Table 6. No significant difference was observed between the blanched dried and microwaved cooked samples (BLDR, BLDO, and FLMCO2), indicating a moderate reduction of phytates across all methods. Furthermore, the absence of a significant difference between the unprocessed fresh leaves and the un-blanched leaves dried at room temperature may be attributed to Shade drying entails limited processing, thereby retaining certain compounds similar to those found in fresh leaves. The lowest total tannin content was observed in the fermented and ten-minute cooked samples, measuring 40.28 mg/100 g and 17.8077 mg/100 g, respectively. In spontaneous fermentation, phytase enzymes generated by fermenting microorganisms hydrolyze phytic acid, the storage form of phosphorus in plants, into inositol and free phosphate, resulting in a significant reduction of phytate levels in FFL samples (Rollán et al., 2019). Naseem et al. (2023) reported a significant reduction of phytates in fermented spinach. Furthermore, research conducted by Irakoze et al. (2023) and Sangija et al. (2022) indicated a decrease in phytates in fermented nightshade species, corroborating the results of the present study. Several studies indicate that blanching and boiling vegetables decrease the phytate content in ILVs (Ngungulu et al., 2024; Mwanri et al., 2018; Tiisekwa et al., 2024), which is consistent with the findings of the current study. Phytates bind essential minerals, forming insoluble complexes that decrease their bioavailability (Aderibigbe et al., 2022). Phytic acid consumption (4-9 mg/100 g) has been shown to decrease iron absorption by 4 to 5 times (Loukou et al., 2019).

Oxalates: The concentration of oxalates in unprocessed leaves of *Justicia heterocarpa* exhibited significant variation ($P<0.05$) across different processing methods, as shown in Table 6. The concentrations of oxalates did not significantly vary between the cooked samples (FLMCO2, FLCO10, and FLCO5), indicating a substantial reduction of oxalates in these methods relative to other processing techniques.

The minimum reduction of oxalates was observed in unblanched samples dried directly in the room, consistent with the findings of Natesh et al. (2017), which indicated that direct drying and storage did not significantly alter the reduction of oxalic acid in ILVs. Numerous studies indicate that oxalates can be effectively reduced through boiling and blanching (Ngungulu et al., 2024; Mwanri et al., 2018; Natesh et al., 2017), corroborating the findings of the current research. The total oxalic acid content in fresh leaves of *Justicia heterocarpa* is lower than that found in other green leafy vegetables, including spinach (329.6-2350 mg/100 g FW) and rhubarb (1235 mg/100 g FW) (Salgado et al., 2023). Soybeans have been reported to contain lower levels of oxalates (124-497 mg/100 g FW) (Salgado et al., 2023), compared to the fresh leaves of *Justicia heterocarpa*. This suggests that certain indigenous leafy vegetables (ILVs) may have higher concentrations of anti-nutrients than legumes, which are typically recognized for their anti-nutrient content. The lethal dose of oxalate in adults has been documented to range from 10 to 15 grams per body weight (Ngungulu et al., 2024).

4. CONCLUSION AND RECOMMENDATION

The present study examined how processing affects *Justicia heterocarpa*'s nutritional and anti-nutritional characteristics. Cooking for ten minutes and blanching before drying reduced anti-nutrients more than unblanched samples. Beta-carotene, vitamin C, and minerals were moderately preserved in microwaved cooked and unblanched shade-dried samples. Fermented leaves had the highest Total Phenolic Compounds (TPC), which protect against non-communicable diseases. The findings are important for Tanzanian communities with dietary inadequacies. To promote *Justicia heterocarpa* as a functional food across various dietary patterns, further research is necessary to ascertain its phytochemical composition and antioxidant activity, as well as the role these compounds may play in the prevention of chronic diseases such as cancer and heart disease.

DATA AVAILABILITY

The data produced and examined in this study can be obtained from the corresponding author upon a reasonable request.

FUNDING

The study was funded by the Sokoine University of Agriculture (SUA) under Higher Education for Economic Transformation (HEET) project.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies, such as large language models (ChatGPT, COPILOT, etc.) and text-to-image generators, have been used during the writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Aderibigbe, O. R., Ezekiel, O. O., Owolade, S. O., Korese, J. K., Sturm, B., & Hensel, O. (2022). Exploring the potentials of underutilized grain amaranth (*Amaranthus* spp.) along the value chain for food and nutrition security: A review. *Critical Reviews in Food Science and Nutrition*, 62(3), 656–669.
<https://doi.org/10.1080/10408398.2020>
- Ahmed, A., Almas, S., Kissaka, K., & Suleiman, R. (2023). Effect of sun-dry on nutritional and sensory acceptability of wilted African leafy vegetables: A case study of Morogoro Region, Tanzania. *Frontiers in Sustainable Food Systems*, 7(June), 1–9.
- Alassane, C. T., Abdoulaye, T., Fabrice, Z. A., Ahmont, K., & Claude, L. (2022). Effect of three drying modes on nutritive and antinutritive properties of leafy vegetables consumed in northern Côte d'Ivoire. *1873*(4), 102–111.
- Atuna, R. A., Djah, J., Achaglinkame, M. A., Bakker, S., Dari, L., Osei-Kwarteng, M., et al. (2022). Types of indigenous vegetables consumed, preparation, preferences, and perceived benefits in Ghana. *Journal of Ethnic Foods*, 9(1).
<https://doi.org/10.1186/s42779-022-001>
- Bighaghire, R., Okidi, L., Muggaga, C., & Ongeng, D. (2021). Traditional vegetable preservation technologies practiced in the Acholi subregion of Uganda improve mineral bioavailability but negatively affect the contribution of vegetables to household

- needs for micronutrients. *Food Science and Nutrition*, 9(2), 589–604.
- Bokelmann, W., Huyskens-Keil, S., Ferenczi, Z., & Stöber, S. (2022). The role of indigenous vegetables to improve food and nutrition security: Experiences from the project HORTINLEA in Kenya (2014–2018). *Frontiers in Sustainable Food Systems*, 6(April), 1–19.
- Borshchevskaya, L. N., Gordeeva, T. L., Kalinina, A. N., & Sineokii, S. P. (2016). Spectrophotometric determination of lactic acid. *Journal of Analytical Chemistry*, 71(8), 755–758.
- Bukenya, R., Laborde, J. E. A., Mamiro, P., Mugabi, R., & Kinabo, J. (2023). Assessment of nutrient adequacy of complementary foods for infants and young children in Morogoro, Tanzania. *Scientific African*, 19.
- Chacha, J. S., & Laswai, H. S. (2020). Traditional practices and consumer habits regarding consumption of underutilized vegetables in Kilimanjaro and Morogoro regions, Tanzania. *International Journal of Food Science*, 2020.
- Degrain, A., Manhivi, V., Remize, F., Garcia, C., & Sivakumar, D. (2020). Effect of lactic acid fermentation on color, phenolic compounds, and antioxidant activity in African nightshade. *Microorganisms*, 8(9), 1–12.
- Duguma, H. T. (2020). Wild edible plant nutritional contribution and consumer perception in Ethiopia. *International Journal of Food Science*, 2020.
- Farag, M. A., Abib, B., Qin, Z., Ze, X., & Ali, S. E. (2023). Dietary macrominerals: Updated review of their role and orchestration in human nutrition throughout the life cycle with sex differences. *Current Research in Food Science*, 6(January), 100254.
- Fombang, E. N., & Mbofung, C. M. F. M. (2017). The effect of steam blanching and drying method on nutrients, phytochemicals, and antioxidant activity of *Moringa (Moringa oleifera L.)* leaves. *American Journal of Food Science and Technology*, 5(2), 53–60. <http://pubs.sciepub.com/ajfst/5/2/4>
- Gowele, V. F., Kinabo, J., Jumbe, T., Kirschmann, C., Frank, J., & Stuetz, W. (2019). Provitamin A carotenoids, tocopherols, ascorbic acid, and minerals in indigenous leafy vegetables from Tanzania. *Foods*, 8(1), 7–10.
- Heaney, D., Padilla-Zakour, O. I., & Chen, C. (2024). Processing and preservation technologies to enhance indigenous food sovereignty, nutrition security, and health equity in North America. *Frontiers in Nutrition*, 11(June), 1–8.
- Henze, J., Abukutsa-Onyango, M., & Opiyo, A. M. (2020). Production and marketing of African indigenous leafy vegetables. *Training Manual for Extension Officers and Practitioners*. https://edoc.huberlin.de/bitstream/handle/18452/23783/Hortinlea2020_Production_and_Marketing_of_African_Indigenous-Leafy_Vegetables.pdf?sequence=1
- Imathiu, S. (2021). Neglected and underutilized cultivated crops with respect to indigenous African leafy vegetables for food and nutrition security. 9(3), 115–125.
- Irakoze, M. L., Owaga, E. E., & Wafula, E. N. (2023). Effect of lactic acid bacteria fermentation on nutrients and anti-nutrients of African black nightshade and African spider plant. *Scientific African*, 21, e01762. <https://doi.org/10.1016/j.sciaf.2023.e01762>
- Irakoze, M. L., Wafula, E. N., & Owaga, E. (2021). Potential role of African fermented indigenous vegetables in maternal and child nutrition in Sub-Saharan Africa. *International Journal of Food Science*, 2021(SDG 2).
- Issa-Zacharia, A., Majaliwa, N. K., Nyamete, F. A., & Chove, L. M. (2024). Diversity of underutilized vegetables in Africa and their potential in the reduction of micronutrient deficiency: A review. *World Journal of Food Science and Technology*, 8(1), 1–13.
- Jachimowicz, K., Winiarska-Mieczan, A., Baranowska-Wójcik, E., & Bąkowski, M. (2021). Pasta as a source of minerals in the diets of Poles; effect of culinary processing of pasta on the content of minerals. *Foods*, 10(9), 1–18.
- Jeje, O. A., O, G. I., & OT, O. A. (2023). Mitigation of hidden hunger through biofortification: An appraisal. *International Journal of Recent Innovations in Food Science and Nutrition*, 6(1), 13–29.
- John, S. E., Azizi, K., Hancy, A., Twin'omujuni, A., Katana, D., Shine, J., et al. (2023). The prevalence and risk factors associated with iron, vitamin B12, and folate deficiencies in pregnant women: A cross-sectional study in Mbeya, Tanzania. *PLOS Global Public Health*, 3(4), 1–15.

- <https://doi.org/10.1371/journal.pgph.0001828>
- Kandongga, A. (2019). Effect of blanching and drying on micronutrients and anti-nutritional factors in false sesame and common bean leaves. *Master's thesis*, Nelson Mandela African Institution of Science and Technology.
- Khasanah, N. U., & Riyanto, R. (2023). Analysis of ascorbic acid content in pineapples and melons using the UV-Vis spectrophotometry method. *Indonesian Journal of Applied Science and Engineering*, 1(1), 1–6.
- Kimura, M., & Rodriguez-Amaya, D. B. (2002). A scheme for obtaining standards and HPLC quantification of leafy vegetable carotenoids. *Food Chemistry*, 78(3), 389–398.
- Kiringia, D. (2019). African leafy vegetables pre-harvest and post-harvest constraints and. *ResearchGate*.
<https://www.researchgate.net/publication/325323882>
- Leonard, W., Zhang, P., Ying, D., Adhikari, B., & Fang, Z. (2021). Fermentation transforms the phenolic profiles and bioactivities of plant-based foods. *Biotechnology Advances*, 49, 107763.
<https://doi.org/10.1016/j.biotechadv.2021.107763>
- Li, X., Yadav, R., & Siddique, K. H. M. (2020). Neglected and underutilized crop species: The key to improving dietary diversity and fighting hunger and malnutrition in Asia and the Pacific. *Frontiers in Nutrition*, 7(November), 1–12.
- Loukou, A. L., Anvoh, K. Y. B., Kouakou, K. H., & Brou, K. (2019). Nutritional composition and bioavailability prediction of calcium, iron, zinc, and magnesium in *Justicia galeopsis* leaves in Côte d'Ivoire. *International Journal of Biological and Chemical Sciences*, 12(6), 2615–2623.
- Martins, T., Barros, A. N., Rosa, E., & Antunes, L. (2023). Enhancing health benefits through chlorophylls and chlorophyll-rich agro-food: A comprehensive review. *Molecules*, 28(14).
- Mieszczakowska-Frać, M., Celejewska, K., & Płocharski, W. (2021). Impact of innovative technologies on the content of vitamin C and its bioavailability from processed fruit and vegetable products. *Antioxidants*, 10(1), 1–19.
- Misci, C., Taskin, E., Dall'Asta, M., Fontanella, M. C., Bandini, F., Imathiu, S., et al. (2021). Fermentation as a tool for increasing food security and nutritional quality of indigenous African leafy vegetables: The case of *Cucurbita* sp. *Food Microbiology*, 99 (December 2020), 103820.
<https://doi.org/10.1016/j.fm.2021.103820>
- Mohankumar, J. B., Uthira, L., & Su, M. (2018). Total phenolic content of organic and conventional green leafy vegetables. 2(1).
- Mungofa, N., Sibanyoni, J. J., Mashau, M. E., & Beswa, D. (2022). Prospective role of indigenous leafy vegetables as functional food ingredients. *Molecules*, 27(22), 1–23.
- Mwanri, A. W., Mamboleo, T. F., Msuya, J. M., & Goweke, V. F. (2018). Oxalate, phytate, and nitrate content in African nightshade, spider plant, and amaranths at different stages of maturity. *African Journal of Food Science*, 12(11), 316–322.
- Naseem, A., Akhtar, S., Ismail, T., Qamar, M., Sattar, D. E. S., Saeed, W., et al. (2023). Effect of growth stages and lactic acid fermentation on anti-nutrients and nutritional attributes of spinach (*Spinacia oleracea*). *Microorganisms*, 11(9), 1–15.
- Natesh, N. H., Akhtar, S. K., & A, L. A. (2017). An overview of nutritional and anti-nutritional factors in green leafy vegetables. *Horticultural International Journal*, 1(2), 58–65.
- Ngungulu, T. P., Wenaty, A., Chove, B., Suleiman, R., & Mbwana, H. (2024). Effect of different processing methods on the anti-nutritive components of selected indigenous vegetables. *Asian Food Science Journal*, 23(8), 80–94.
- Nicholaus, C., Martin, H. D., Kassim, N., Matemu, A. O., & Kimiywe, J. (2020). Dietary practices, nutrient adequacy, and nutrition status among adolescents in boarding high schools in the Kilimanjaro region, Tanzania. *Journal of Nutrition and Metabolism*, 2020.
- Ogbonnaya, F. C., Ukangwa, N. A., Amah, G. H., & Oladele, L. O. (2024). Inductively coupled plasma optical electro spectroscopic characterization of some indigenous Nigerian leafy vegetables as prospective mineral sources in human

- diets. *Nigerian Journal of Pure and Applied Sciences*, 37(1), 4772–4779.
- Oluoch, M. O., & Habwe, F. (2012). Food preparation and processing methods on nutrient retention and accessibility in selected indigenous vegetables from East Africa promote sustainable agriculture in Borno State. View the project vegetable breeding and seed systems for poverty reduction in A. *ResearchGate*. <https://www.researchgate.net/publication/316990800>
- Piyarach, K., Nipawan, K., Chadapon, C., Daluwan, S., & Kunjana, R. (2020). Effect of drying on β -carotene, α -carotene, lutein, and zeaxanthin content in vegetables and its application for vegetable seasoning. *E3S Web of Conferences*, 141, 1–6.
- Rollán, G. C., Gerez, C. L., & Leblanc, J. G. (2019). Lactic fermentation as a strategy to improve the nutritional and functional values of pseudocereals. *Frontiers in Nutrition*, 6(July), 1–9.
- Salgado, N., Silva, M. A., Figueira, M. E., Costa, H. S., & Albuquerque, T. G. (2023). Oxalate in foods: Extraction conditions, analytical methods, occurrence, and health implications. *Foods*, 12(17).
- Sánchez-Rangel, J. C., Benavides, J., Heredia, J. B., Cisneros-Zevallos, L., & Jacobo-Velázquez, D. A. (2013). The Folin-Ciocalteu assay revisited: Improvement of its specificity for total phenolic content determination. *Analytical Methods*, 5(21), 5990–5999.
- Sangija, F., Kazosi, M., Martin, M., & Matemu, A. (2022). Trends and constraints in the utilization of African nightshade (*Solanum nigrum* complex) in Tanzania: A case study of Kilimanjaro and Morogoro regions. *African Journal of Food, Agriculture, Nutrition, and Development*, 22(6), 20623–20645.
- Sangija, F., Martin, H., & Matemu, A. (2022). Effect of lactic acid fermentation on the nutritional quality and consumer acceptability of African nightshade. *Food Science and Nutrition*, 10(9), 3128–3142.
- Sarfo, J., Pawelzik, E., & Keding, G. B. (2022). Are processed fruits and vegetables able to reduce diet costs and address micronutrient deficiencies? Evidence from rural Tanzania. *Public Health Nutrition*, 25(9), 2637–2650.
- Schoofs, H., Schmit, J., & Rink, L. (2024). Zinc toxicity: Understanding the limits. *Molecules*, 29(13).
- Stevens, G. A., Beal, T., Mbuya, M. N. N., Luo, H., Neufeld, L. M., Addo, O. Y., et al. (2022). Micronutrient deficiencies among preschool-aged children and women of reproductive age worldwide: A pooled analysis of individual-level data from population-representative surveys. *Lancet Global Health*, 10(11), e1590–e1599.
- Stoll, D. A., Wafula, E. N., Mathara, J. M., Trierweiler, B., Kulling, S. E., & Huch, M. (2021). Fermentation of African nightshade leaves with lactic acid bacterial starter cultures. *International Journal of Food Microbiology*, 342 (January 2020), 109056. <https://doi.org/10.1016/j.ijfoodmicro.2021.109056>
- Stuetz, W., Gowele, V., Kinabo, J., Bundala, N., Mbwana, H., Rybak, C., et al. (2019). Consumption of dark green leafy vegetables predicts vitamin A and iron intake and status among female small-scale farmers in Tanzania. *Nutrients*, 11(5), 1–18.
- Tanor, E. B., Ntlatlapa, T., & Magama, S. (2020). Assessment of the nutritional value of selected wild leafy vegetables growing in the Roma Valley, Lesotho. *European Journal of Nutrition and Food Safety*, 12(10), 32–45.
- Tiisekwa, A. B., Chove, B. E., Mongi, R., Rybak, C., & Stuetz, W. (2024). Carotenoids, tocopherols, minerals, and phytate in fresh and sun-dried dark green leafy vegetable varieties from the Lindi region, Tanzania. *Technologies in Horticulture*, 4(1), 0–0.
- Traoré, K., Parkouda, C., Savadogo, A., Ba/Hama, F., Kamga, R., & Traoré, Y. (2017). Effect of processing methods on the nutritional content of three traditional vegetables leaves: Amaranth, black nightshade, and jute mallow. *Food Science and Nutrition*, 5(6), 1139–1144.
- Ukom, A., Albert, M., Ojimekwe, P., Offia-Olua, B., & Nwanagba, L. (2023). Impact of cooking methods on the chemical and antioxidant composition of some indigenous vegetables used in different food dishes in Southeast Nigeria. *Journal of Ethnic Foods*, 10(1). <https://doi.org/10.1186/s42779-023-00110-0>
- Verardo, V., Gómez-Caravaca, A. M., & Tabanelli, G. (2020). Bioactive components in fermented foods and food by-products. *Foods*, 9(2), 4–6.

Zong, C., Wu, Q., Wu, A., Chen, S., Dong, D., Zhao, J., et al. (2021). Exploring the diversity mechanism of fatty acids and the loss mechanisms of polyunsaturated fatty acids and fat-soluble vitamins in alfalfa silage using different additives. *Animal Feed Science and Technology*, 280(January), 115044.
<https://doi.org/10.1016/j.anifeedsci.2021.115044>

Disclaimer/Publisher's Note: The statements, opinions, and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions, or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://pr.sdiarticle5.com/review-history/130549>